

NOTES ON CARTIER-DIEUDONNÉ THEORY

CHING-LI CHAI

version 0.01

The purpose of this set of notes is to serve as an introduction to the Cartier-Dieudonné theory on commutative smooth formal groups. The upshot of the Cartier theory is to transform most of the questions about commutative formal groups to questions in linear algebra: The main theorem 3.3 says that the category of smooth commutative formal groups over a commutative ring k is equivalent to a suitable full subcategory of the category of left modules over a non-commutative ring $\text{Cart}(k)$. The equivalence above is a sort of Morita equivalence. When the ring k is a $\mathbb{Z}_{(p)}$ -algebra, where p is a prime number, there is a local version of the main theorem, with the ring $\text{Cart}(k)$ replaced by a subring $\text{Cart}_p(k)$ of $\text{Cart}(k)$ defined by an idempotent in $\text{Cart}(k)$.

A key role is played by a smooth commutative formal group Λ , which is the formal completion of a restricted version of the group scheme of universal Witt vectors; see 1.6 for its definition. This smooth formal group Λ is in some sense a free generator of the additive category of smooth commutative formal groups. The ring $\text{Cart}(k)$ is the opposite ring of $\text{End}_k(\Lambda)$; it is in natural bijection with the set of all formal curves in Λ .

The best textbook of Cartier theory is the elegant booklet [Z] by Zink, where the exposition in §2 of [R] is fully developed. Lazard's monograph [L] is the first complete documentation of Cartier's theory, but may be hard to digest because of its style. Hazewinkel's treatment [H] employs the technology of the "functional equation lemma" and does everything two or three times; it is a very useful reference with 573+ix pages and a good indexing system.

Although the main results of Cartier theory does not depend on the Witt vectors, in applications the Witt vectors are often very useful. The basic properties of both the ring of universal Witt vectors and the ring of p -adic Witt vectors can be found in Appendix A; the exposition there is independent of the rest of this set of notes. The Witt vectors can also be viewed as part of the Cartier theory, for they are the Cartier module attached to the formal completion $\widehat{\mathbb{G}}_m$ of \mathbb{G}_m in the two versions of Cartier theory. The universal Witt vectors are formal curves in $\widehat{\mathbb{G}}_m$, and the p -adic Witt vectors are the p -typical formal curves in $\widehat{\mathbb{G}}_m$.

§1. Formal groups

In this section k denotes a commutative ring with 1. The notion of smooth formal groups adopted here is slightly different from the standard definition, because we consider them as functors on the category of nilpotent algebras.

(1.1) Definition Let \mathfrak{Nil}_k be the category of all nilpotent k -algebras, consisting of all commutative k -algebras N without unit such that $N^n = 0$ for some positive integer n .

(1.1.1) Remark Clearly \mathfrak{Nilp}_k is isomorphic to the category of augmented k -algebras $k \rightarrow R \xrightarrow{\epsilon} k$ such that the augmentation ideal $I = \text{Ker}(\epsilon)$ is nilpotent; the isomorphisms are given by $N \mapsto k \oplus N$ and $(R, \epsilon) \mapsto \text{Ker}(\epsilon)$.

(1.1.2) Definition Let $\mathfrak{ProNilp}_k$ be the category of all filtered projective limits of nilpotent k -algebras. Every functor $G : \mathfrak{Nilp}_k \rightarrow \mathfrak{Sets}$ can be uniquely extended to a functor from \mathfrak{Nilp} to \mathfrak{Sets} which commutes with filtered projective limits; this extension is also denoted by G . The analogous statement holds for functors from \mathfrak{Nilp}_k to \mathfrak{Ab} .

Remark As an example, let $k[[X_1, \dots, X_n]]$ be a power series ring over k in n variables. Denote by $k[[X_1, \dots, X_n]]^+$ the subset of $k[[X_1, \dots, X_n]]$ consisting of all power series whose constant term is 0. Then $k[[X_1, \dots, X_n]]^+$ is an object in $\mathfrak{ProNilp}_k$, and

$$G(k[[X_1, \dots, X_n]]^+) = \varprojlim_{i \geq 1} G(k[[X_1, \dots, X_n]]^+ / ((X_1, \dots, X_n)k[[X_1, \dots, X_n]]^i) .$$

(1.2) Definition Let $G : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be a functor from \mathfrak{Nilp}_k to the category of all abelian groups.

- (1) The functor G is *left exact* if it commutes with finite inverse limits and $G(0) = (0)$. (Actually the latter condition is a special case of the first one: take the inverse limit over the empty indexing set.)
- (2) The functor G is *formally smooth* if every surjection $N_1 \rightarrow N_2$ in \mathfrak{Nilp}_k induces a surjection $G(N_1) \rightarrow G(N_2)$.
- (3) The functor G is *right exact* if it commutes with finite direct product, and every exact sequence $N_3 \rightarrow N_2 \rightarrow N_1 \rightarrow 0$ in \mathfrak{Nilp}_k induces an exact sequence $G(N_3) \rightarrow G(N_2) \rightarrow G(N_1) \rightarrow 0$ in \mathfrak{Ab} .
- (4) The functor G is *weakly left exact* if G commutes with finite direct product, and if for every short exact sequence $0 \rightarrow N_1 \rightarrow N_2 \xrightarrow{\pi} N_3 \rightarrow 0$ in \mathfrak{Nilp}_k such that $N_3^2 = (0)$ and N_3 is a free k -module, the induced sequence $0 \rightarrow G(N_1) \rightarrow G(N_2) \rightarrow G(N_3) \rightarrow 0$ is short exact.
- (5) The functor G is *half exact* if G commutes with finite direct product, and if for every exact sequence $0 \rightarrow N_1 \rightarrow N \xrightarrow{\pi} N_2 \rightarrow 0$ in \mathfrak{Nilp}_k such that $N_1 \cdot N = (0)$, the group $G(N_1)$ operates simply transitively on $N(\pi)^{-1}(\xi)$ for every $\xi \in G(N_2)$ such that $N(\pi)^{-1}(\xi) \neq \emptyset$.

Remark Left exactness implies weak left exactness.

(1.2.1) Definition Let k be a commutative ring with 1 and let \mathfrak{Mod}_k be the category of k -modules. There is a natural embedding of \mathfrak{Mod}_k into \mathfrak{Nilp}_k , endowing each k -module M the trivial multiplication structure, i.e. $M \cdot M = (0)$. Let G be a functor from \mathfrak{Nilp}_k to \mathfrak{Ab}

which commutes with finite direct sums. The *tangent functor* $t_G : \mathfrak{Mod}_k \rightarrow \mathfrak{Mod}_k$ of G is defined by restricting G to \mathfrak{Mod}_k and endowing $G(M)$ the natural k -module structure for any k -module M .

(1.3) Definition A functor $G : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ from \mathfrak{Nilp}_k to the category of abelian groups is a *smooth commutative formal group* if G is left exact, formally smooth, and commutes with arbitrary direct sums. Since the formal groups we will be concerned mostly with are smooth and commutative, we will call them *smooth formal groups* for short unless specified otherwise.

(1.3.1) Definition Let k be a commutative ring with 1 and let I be an indexing set.

- (i) Let $\underline{X} = (X_i)_{i \in I}$ be a set of variables indexed by the set I . Denote by $k[[\underline{X}]] = k[[X_i]]_{i \in I}$ the inverse limit of all formal power series rings $k[[X_j]]_{j \in J}$ where J runs through all finite subsets of I . In other words, $k[[\underline{X}]] = k[[X^I]]$ consists of all formal power series

$$\sum_{\alpha} a_{\alpha} X_{\alpha}^{n_{\alpha}}, \quad a_{\alpha} \in k,$$

where α runs through all functions $\alpha : I \rightarrow \mathbb{N}$ vanishing outside some finite subset of I . Elements of $k[[\underline{X}]] = k[[X^I]]$ are in bijection with k -valued functions on the set of all monomials in the variables \underline{X} .

- (ii) Denote by $k[[\underline{X}]]^+$ the augmentation ideal of $k[[\underline{X}]]$, consisting of all power series without the constant term. For each $n \geq 1$, the quotient $k[[\underline{X}]]^+ / (k[[\underline{X}]]^+)^n$ is a nilpotent k -algebra, and $k[[\underline{X}]]^+$ is the filtered inverse limit of the $k[[\underline{X}]]^+ / (k[[\underline{X}]]^+)^n$'s.
- (iii) Denote by $\widehat{\mathbb{A}}^{(I)}$ the functor from \mathfrak{Nilp}_k to \mathfrak{Sets} such that

$$\widehat{\mathbb{A}}^{(I)}(N) = \bigoplus_{i \in I} N,$$

the set underlying the direct sum of I copies of N . Clearly elements of $k[[X^{(I)}]]$ gives rise to formal functions on $\widehat{\mathbb{A}}^{(I)}$, i.e. maps from $\widehat{\mathbb{A}}^{(I)}$ to $\widehat{\mathbb{A}}^1$.

(1.3.2) Definition Let k be a commutative ring with 1 and let I be a set. A commutative formal group law on $\widehat{\mathbb{A}}^{(I)}$ is morphism $\mu : \widehat{\mathbb{A}}^{(I)} \times \widehat{\mathbb{A}}^{(I)} \rightarrow \widehat{\mathbb{A}}^{(I)}$ which provides a commutative group law on $\widehat{\mathbb{A}}^{(I)}$. Equivalently, a commutative formal group law is a homomorphism $\mu^* : k[[X^{(I)}]] \rightarrow k[[X^{(I)}, Y^{(J)}]]$ which is coassociative, cocommutative, and admits a coinverse. Often we identify μ^* with its restriction to the free topological generators \underline{X} .

(1.4) Some examples of smooth formal groups.

(1.4.1) The formal group $\widehat{\mathbb{G}}_a$ attached to the additive group:

$$\widehat{\mathbb{G}}_a(N) := N,$$

the additive group underlying the nilpotent k -algebra N .

(1.4.2) The formal group $\widehat{\mathbb{G}}_m$ attached to the multiplicative group:

$$\widehat{\mathbb{G}}_m(N) := 1 + N \subset (k \oplus N)^\times \quad \forall N \in \mathfrak{Nilp}_k.$$

Here $(k \oplus N)^\times$ denotes the group of units of the augmented k -algebra $k \oplus N$, so the group law is $(1 + u) \cdot (1 + v) = 1 + u + v + uv$ for $u, v \in N$.

(1.5) The Lubin-Tate formal group.

(1.5.1) Let \mathcal{O} be a complete discrete valuation ring whose residue field κ is a finite field with $q = p^a$ elements, where p is a prime number. Let π be a uniformizing element of \mathcal{O} . Recall that a *Lubin-Tate* formal group over \mathcal{O} is a one-dimensional smooth formal group $G = \mathrm{Spf}(\mathcal{O}[[x]])$ over \mathcal{O} with an endomorphism $\phi : G \rightarrow G$ such that

$$\phi(x) := \phi^*(x) \equiv \pi x + x^q \pmod{(\pi, x^2)}.$$

It is known that every polynomial $\phi(x) \in \mathcal{O}[[x]]$ satisfying the above property uniquely determines a formal group law $\Phi_\phi(x, y)$ on $\mathrm{Spf}(\mathcal{O}[[x]])$ such that $\phi(x)$ defines an endomorphism of $\Phi_\phi(x, y)$, and there is a ring homomorphism $\mathcal{O} \rightarrow \mathrm{End}(\Phi_\phi)$ such that π operates on the formal group law $\Phi_\phi(x, y)$ via $\phi(x)$. Moreover for any two Lubin-Tate formal groups $\Phi_{\phi_1}, \Phi_{\phi_2}$ over \mathcal{O} there exists a unique \mathcal{O} -equivariant isomorphism $\psi : \Phi_{\phi_1} \xrightarrow{\sim} \Phi_{\phi_2}$ such that $\psi(x) \equiv x \pmod{x^2}$.

(1.5.2) Let \mathcal{O}, π be as above, $q = \mathrm{Card}(\mathcal{O}/\pi\mathcal{O})$. Let K be the fraction field of \mathcal{O} . Let

$$f_\pi(X) = \sum_{n \geq 0} \frac{X^{q^n}}{\pi^n} = X + \frac{X^q}{\pi} + \frac{X^{q^2}}{\pi^2} + \cdots \in K[[X]].$$

Let $\Phi_\pi(X, Y) = f_\pi^{-1}(f_\pi(X) + f_\pi(Y))$. A priori $\Phi_\pi(X, Y)$ has coefficients in K , but in fact $\Phi_\pi(X, Y) \in \mathcal{O}[[X, Y]]$. This can be proved directly, or one can use the “functional equation lemma” on p.9 of [H], since $f_\pi(X)$ satisfies the functional equation

$$f_\pi(X) = X + \frac{1}{\pi} f_\pi(X^q).$$

It follows that $\Phi_\pi(X, Y)$ is a one-dimensional formal group law, and $f_\pi(X)$ is the logarithm of $\Phi_\pi(X, Y)$. Moreover one checks that the polynomial

$$\phi_\pi(X) := f_\pi(\pi f_\pi(X))$$

has coefficients in \mathcal{O} and satisfies

$$\phi_\pi(X) \equiv \pi X + X^q \pmod{\pi x^2 \mathcal{O}[[x]]}.$$

Hence $\Phi_\pi(x, y)$ is a Lubin-Tate formal group law for (\mathcal{O}, π) .

(1.6) Definition We define a restricted version of the smooth formal group attached to the universal Witt vector group, denoted by Λ :

$$\Lambda(N) = 1 + t k[t] \otimes_k N \subset ((k \oplus N)[t])^\times \quad \forall N \in \mathfrak{Nilp}_k.$$

In other words, the elements of $\Lambda(N)$ consists of all polynomials of the form $1 + u_1 t + u_2 t^2 + \dots + u_r t^r$ for some $r \geq 0$, where $u_i \in N$ for $i = 1, \dots, r$. The group law of $\Lambda(N)$ comes from multiplication in the polynomial ring $(k \oplus N)[t]$ in one variable t . The formal group Λ will play the role of a free generator in the category of (smooth) formal groups. When we want to emphasize that the polynomial $1 + \sum_{i \geq 1} u_i t^i$ is regarded as an element of $\Lambda(N)$, we denote it by $\lambda(1 + \sum_{i \geq 1} u_i t^i)$.

(1.6.1) Remark (i) $\Lambda(k[[X]]^+)$ consists of all formal power series in $k[[X, t]]$ of the form

$$1 + \sum_{m, n \geq 1} b_{m, n} X^m t^n, \quad b_{m, n} \in k$$

such that for every m , there exists an integer $C(m)$ such that $b_{m, n} = 0$ for all $n \geq C(m)$.

(ii) The formal completion \widetilde{W}^\wedge of the universal Witt vector group \widetilde{W} , defined in §A, is given by

$$\widetilde{W}^\wedge(N) = 1 + t N[[t]] \subset ((k \oplus N)[[t]])^\times \quad \forall N \in \mathfrak{Nilp}_k.$$

In particular $\widetilde{W}^\wedge(k[[X]]^+)$ consists of *all* power series $1 + \sum_{m, n \geq 1} b_{m, n} X^m t^n$, $b_{m, n} \in k$ in $k[[X, t]]$.

(1.6.2) Exercise Prove that for every nilpotent k -algebra N , every element of $\Lambda(N)$ can be uniquely expressed as a finite product

$$(1 - a_1 t)(1 - a_2 t^2) \cdots (1 - a_m t^m)$$

with $a_1, \dots, a_m \in N$. Deduce that

$$\Lambda(k[[X]]^+) = \left\{ \prod_{m, n \geq 1} (1 - a_{mn} X^m t^n) \mid a_{m, n} \in k, \forall m \exists C_m > 0 \text{ s.t. } a_{mn} = 0 \text{ if } n \geq C_m \right\}$$

§2. The Cartier ring

(2.1) **Definition** Let k be a commutative ring with 1. Let $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be a functor from the category of commutative nilpotent k -algebras to the category of abelian groups, extended to the category of topologically nilpotent k -algebras by filtered inverse limit as in 1.1.2. We say that H is *weakly symmetric*, or equivalently that H satisfies the *weak symmetry condition*, if for every $n \geq 1$, the natural map

$$H((k[[T_1, \dots, T_n]]^+)^{S_n}) \rightarrow H(k[[T_1, \dots, T_n]]^+)^{S_n}$$

induced by the inclusion $k[[T_1, \dots, T_n]]^{S_n} \hookrightarrow k[[T_1, \dots, T_n]]$ is an isomorphism. Here S_n is the symmetric group in n letters operating naturally on the power series ring $k[[T_1, \dots, T_n]]$. Note that $k[[T_1, \dots, T_n]]^{S_n}$ is the power series ring generated by the elementary symmetric polynomials in the variables T_1, \dots, T_n .

(2.1.1) **Lemma** *Let k be a commutative ring with 1. Let $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be a functor.*

- (i) *Suppose that H is left exact, that is H commutes with finite inverse limits. Then H is weakly symmetric. In particular this is the case if H is a smooth commutative formal group over k .*
- (ii) *If H is weakly left exact, then H is weakly symmetric.*

PROOF. The ring $k[[T_1, \dots, T_n]]^{S_n}$ is a fiber product of two ring homomorphisms from $k[[T_1, \dots, T_n]]$ to $\prod_{\sigma \in S_n} k[[T_1, \dots, T_n]]$; one is the diagonal embedding, the other sends $f(\underline{T})$ to $(f(\underline{T}^\sigma))_{\sigma \in S_n}$. Applying the half-exactness of H to this fiber product, one deduces (i). The stronger statement (ii) follows from the same argument. ■

(2.2) **Theorem** *Notation as in (2.1), and assume that $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ satisfies the weak symmetry condition. Let Λ be the functor defined in (1.6). Then the map*

$$Y_H : \text{Hom}(\Lambda_k, H) \rightarrow H(k[[X]]^+)$$

which sends each homomorphism $\alpha : \Lambda \rightarrow H$ of group-valued functors to the element $\alpha_{k[[X]]^+}(1 - Xt) \in H(k[[X]]^+)$ is a bijection.

Remark (i) Thm. (2.2) can be regarded as a sort of Yoneda isomorphism. The inverse of Y_H is given in the proof.

- (ii) The formal group Λ is in some sense a free generator of the additive category of smooth commutative formal groups, a phenomenon reflected in Thm. 2.2.

PROOF. Suppose that $\alpha \in \text{Hom}(\Lambda, H)$. Given any nilpotent k -algebra N and any element $f(t) = 1 + u_1 t + u_2 t^2 + \dots + u_n t^n \in \Lambda(N)$, we explain why the element $\alpha_N(f) \in H(N)$ is determined by the element $h_\alpha := \alpha_{k[[X]]^+}(1 - Xt) \in H(k[[X]]^+)$.

Let $\beta = \beta_{f,n} : k[[U_1, \dots, U_n]]^+ \rightarrow N$ be the continuous k -linear homomorphism such that $\beta(U_i) = (-1)^i u_i$. Let $\delta = \delta_n : k[[U_1, \dots, U_n]]^+ \rightarrow k[[X_1, \dots, X_n]]^+$ be the continuous homomorphism sending each U_i to the i -th elementary polynomial in the variables X_1, \dots, X_n . Clearly

$$\alpha_N(f) = H(\beta)\alpha_{k[[U_1, \dots, U_n]]^+}(1 - U_1 t + \dots + (-1)^n U_n t^n).$$

Moreover we have

$$\alpha_{k[[U_1, \dots, U_n]]^+}(1 - U_1 t + \dots + (-1)^n U_n t^n) = \sum_{i=1}^n \alpha_{k[[X_1, \dots, X_n]]}(1 - X_i t) = \sum_{i=1}^n H(\iota_i)(h_\alpha)$$

in $H(k[[X_1, \dots, X_n]]^+)^{S_n} = H(k[[U_1, \dots, U_n]]^+) \subset H(k[[X_1, \dots, X_n]]^+)$, where $\iota_i : k[[X]]^+ \rightarrow k[[X_1, \dots, X_n]]^+$ is the continuous k -algebra homomorphism sending X to X_i . The two displayed formulas shows how to compute $\alpha_N(f)$ for any element $f(t) \in H(N)$ in terms of $\alpha_{k[[X]]}(1 - Xt)$. The injectivity of Y_H follows.

Conversely, given an element $h \in H(k[[X]]^+)$, we have to construct a homomorphism of functors $\alpha \in \text{Hom}(\Lambda, H)$ such that $\alpha_{k[[X]]}(1 - Xt) = h$. The argument above provides a procedure to get an element $\alpha_N(f) \in H(N)$ for any element $f(t) \in \Lambda(N)$ for a nilpotent k -algebra N . Explicitly, for $f = 1 + u_1 t + u_2 t^2 + \dots + u_n t^n \in \Lambda(N)$,

- let $\beta_{f,n} : k[[U_1, \dots, U_n]]^+ \rightarrow N$ be the continuous k -linear homomorphism such that $\beta_{f,n}(U_i) = (-1)^i u_i$ for each i ,
- let $j_n : k[[U_1, \dots, U_n]]^+ \hookrightarrow k[[X_1, \dots, X_n]]^+$ be the continuous k -linear injection such that $j(U_i)$ is equal to the i -th elementary symmetric polynomial in X_1, \dots, X_n ,
- let $\iota_i : k[[X]]^+ \rightarrow k[[X_1, \dots, X_n]]^+$ be the continuous k -linear homomorphism such that $\iota_i(X) = X_i$, $i = 1, \dots, n$, and
- let $\tilde{h}_n = \tilde{h}_{f,n} \in H(k[[U_1, \dots, U_n]]^+)$ be the element of $H(k[[U_1, \dots, U_n]]^+)$ such that $H(j_n)(\tilde{h}_n) = \sum_{i=1}^n H(\iota_i)(h)$.

Then $\boxed{\alpha_N(f) = H(\beta_{f,n})(\tilde{h}_{f,n})}$.

It is not hard to check that the element $\alpha_N(f) \in H(N)$ is independent of the choice of the integer n , so that $\alpha_N(f)$ is well-defined. This is left as an exercise, as well as the fact that the collection of maps α_N defines a functor from \mathfrak{Nilp}_k to \mathfrak{Ab} .

Lastly, we verify that $\alpha(f_1 + f_2) = \alpha(f_1) + \alpha(f_2)$ for any $f_1(t), f_2(t) \in H(k[[X]]^+)$. It suffices to check this in the universal case. In other words, it suffices to verify the equality $\alpha(f_1 + f_2) = \alpha(f_1) + \alpha(f_2)$ in $H(k[[U_1, \dots, U_n, V_1, \dots, V_m]]^+)$, where $f_1(t) = 1 - U_1 t + \dots + (-1)^n U_n t^n$ and $f_2(t) = 1 - V_1 t + \dots + (-1)^m V_m t^m$. As above we may assume that U_1, \dots, U_n are the elementary symmetric polynomials in the variables X_1, \dots, X_n and V_1, \dots, V_m are the elementary symmetric polynomials in the variables Y_1, \dots, Y_m . Let ι_i (resp. ι'_j) be the

continuous homomorphism from $k[[X]]$ to $k[[X_1, \dots, X_n, Y_1, \dots, Y_m]]$ such that $\iota_i(X) = X_i$ (resp. $\iota'_j(X) = Y_j$.) Then we have

$$\begin{aligned}\alpha(f_1) &= \sum_{i=1}^n H(\iota_i)(h) \in H((k[[\underline{X}]]^+)^{S_n}) = H((k[[\underline{X}]]^+)^{S_n}) = H(k[[\underline{U}]]^+) \subset H(k[[\underline{U}, \underline{V}]]^+) \\ \alpha(f_2) &= \sum_{j=1}^m H(\iota'_j)(h) \in H((k[[\underline{Y}]]^+)^{S_m}) = H((k[[\underline{Y}]]^+)^{S_m}) = H(k[[\underline{V}]]^+) \subset H(k[[\underline{U}, \underline{V}]]^+)\end{aligned}$$

and

$$\begin{aligned}\alpha(f_1 + f_2) &= \sum_{i=1}^n H(\iota_i)(h) + \sum_{j=1}^m H(\iota'_j)(h) \in H(k[[X_1, \dots, X_n, Y_1, \dots, Y_m]]^+)^{S_{n+m}} \\ &= H((k[[X_1, \dots, X_n, Y_1, \dots, Y_m]]^+)^{S_{n+m}}) \subset H((k[[\underline{X}, \underline{Y}]]^+)^{S_n \times S_m}) = H(k[[\underline{U}, \underline{V}]]^+).\end{aligned}$$

We conclude that $\alpha(f_1 + f_2) = \alpha(f_1) + \alpha(f_2)$. ■

(2.2.1) Corollary *Let $h = h(X, t)$ be an element of $\Lambda(k[[X]]^+)$, and let $\Phi = \Phi_h$ be the endomorphism of Λ_k such that $\Phi_{k[[X]]}(1 - Xt) = h(X, t)$. For each $n \in \mathbb{N}$, define power series ${}^n A_{h,1}(U_1, \dots, U_n), \dots, {}^n A_{h,n}(U_1, \dots, U_n) \in k[[U_1, \dots, U_n]]^+$ by*

$$\prod_{i=1}^n h(X_i, t) = 1 + {}^n A_{h,1}(\sigma_1(\underline{X}), \dots, \sigma_n(\underline{X}))t + \dots + {}^n A_{h,n}(\sigma_1(\underline{X}), \dots, \sigma_n(\underline{X}))t^n,$$

where $\sigma_i(\underline{X})$ denotes the i -th elementary symmetric polynomial in X_1, \dots, X_n .

(i) *Let N be a nilpotent k -algebra, and let $f(t) = 1 + a_1 t + a_2 t^2 + \dots + a_n t^n$ be an element of $\Lambda_k(N)$. Then*

$$\Phi_N(f) = 1 + {}^n A_{h,1}(-u_1, u_2, \dots, (-1)^n u_n) t + \dots + {}^n A_{h,n}(-u_1, u_2, \dots, (-1)^n u_n) t^n.$$

(ii) *We have ${}^{n+1} A_{h,n+1}(U_1, \dots, U_n, 0) = 0$, and*

$${}^{n+1} A_{h,i}(U_1, \dots, U_n, 0) = {}^n A_{h,i}(U_1, \dots, U_n)$$

for each $i = 1, 2, \dots, n$.

(iii) *Suppose that $h(X, t) \equiv 1 \pmod{(X^m)}$, and let $s = \lceil \frac{m}{n} \rceil$. Then*

$${}^n A_{h,i}(U_1, \dots, U_n) \equiv 0 \pmod{(U_1, \dots, U_n)^s}$$

for $i = 1, \dots, n$.

(iv) *In the situation of (i) above, suppose that $N^r = (0)$, then $\Phi_{h,N}(f) = 0$ if $h(X, t) \equiv 1 \pmod{(X^{(r-1)n+1})}$.*

PROOF. The statements (i), (ii) are special cases of Thm. 2.2. The statements (iii), (iv) are easy and left as exercises. ■

(2.3) Definition Define $\text{Cart}(k)$ to be $(\text{End}(\Lambda_k))^{\text{op}}$, the opposite ring of the endomorphism ring of the smooth formal group Λ_k . According to Thm. 2.2, for every weakly symmetric functor $H : \mathfrak{N}ilp_k \rightarrow \mathfrak{A}b$, the abelian group $H(k[[X]]^+) = \text{Hom}(\Lambda_k, H)$ is a *left* module over $\text{Cart}(k)$.

(2.3.1) Definition We define some special elements of the Cartier ring $\text{Cart}(k)$, naturally identified with $\Lambda(k[[X]])$ via $Y = Y_\Lambda$.

$$(i) \quad V_n = Y^{-1}(1 - X^n t), n \geq 1,$$

$$(ii) \quad F_n = Y^{-1}(1 - X t^n), n \geq 1,$$

$$[c] = Y^{-1}(1 - c X t), c \in k.$$

(2.3.2) Lemma For every positive integer n , denote by $\phi_n : k[[X]] \rightarrow k[[X]]$ the k -algebra homomorphism which sends X to X^n . For every $c \in k$, denote by $\psi_c : k[[X]] \rightarrow k[[X]]$ the k -algebra homomorphism which sends X to cX . Then for every weakly symmetric functor $H : \mathfrak{N}ilp \rightarrow \mathfrak{A}b$, we have

$$V_n \gamma = H(\phi_n)(\gamma), \quad [c] \gamma = H(\psi_c)\gamma$$

for every $\gamma \in H(k[[X]])$ and every $c \in k$. Applying the above to Λ , we get

$$V_n [c] F_m = Y^{-1}(1 - c X^n t^m)$$

in $\text{Cart}(k)$.

PROOF. Exercise. ■

(2.3.3) Remark Let \widetilde{W} be the ring scheme of universal Witt vectors defined in §A. For each positive integer n we have endomorphisms V_n, F_n of \widetilde{W} . Consider the element $\omega(1 - X T) \in \widetilde{W}(\mathbb{Z}[[X]])$. Then $V_n(\omega(1 - X T)) = \omega(1 - X T^n)$, and $F_n(\omega(1 - X T)) = \omega(1 - X^n T)$. This contrasts with the notation used in the Cartier ring: $V_n = Y^{-1}(1 - X^n t)$, $F_n = Y^{-1}(1 - X T^n)$, see also Exer. 2.4.2. This kind of "flipping" is inevitable, since $\text{Cart}(\cdot)\mathbb{Z}$ operates on the *right* of \widetilde{W} , and we want the same commutation relation of V_n, F_n with the endomorphisms $[c]$ in 2.4 to hold in all situations.

Remark Often $H(\phi_n)(\gamma)$ is abbreviated as $\gamma(X^n)$, and $H(\psi_c)(\gamma)$ is shortened to $\gamma(cX)$. This is compatible with the standard notation when H is representable as a formal scheme $\text{Spf}R$, where R is an augmented k -algebra complete with respect to the augmentation ideal. The elements of $H(k[[X]])$ are identified with continuous homomorphisms $R \rightarrow k[[X]]$, thought of as "formal curves" in $\text{Spf}R$.

Corollary For every commutative ring with 1 we have

$$\text{Cart}(k) = \left\{ \sum_{m,n \geq 1} V_m [c_{mn}] F_n \mid c_{mn} \in k, \forall m \exists C_m > 0 \text{ s.t. } c_{mn} = 0 \text{ if } n \geq C_m \right\}$$

(2.3.4) Exercise Let k be a commutative ring with 1 and let n be an integer. Prove that n is invertible in $\text{Cart}(k)$ if and only if n is invertible in k .

(2.3.5) Suppose that $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ is weakly symmetric. Let $n \geq 1$ be a positive integer. Denote by $k[\zeta_n]$ the k -algebra $k[T]/(T^n - 1)$, and let $\zeta = \zeta_n$ be the image of T in $k[T]/(T^n - 1)$. Denote by $k[\zeta][[X^{\frac{1}{n}}]]^+$ the $k[\zeta]$ -algebra $k[\zeta][[X, U]]^+ / (U^n - X)k[\zeta][[X, U]]$, and let $X^{\frac{1}{n}}$ be the image of U in $k[\zeta][[X, U]]^+ / (U^n - X)k[\zeta][[X, U]]$. For each $i = 0, \dots, n-1$, let $\phi_{n,i} : k[[X]]^+ \rightarrow k[\zeta][[X^{\frac{1}{n}}]]^+$ be the homomorphism of k -algebras which maps X to $\zeta^i X^{\frac{1}{n}}$. Then

$$F_n \cdot \gamma = \sum_{i=0}^{n-1} H(\phi_{n,i})(\gamma)$$

for every $\gamma \in H(k[[X]]^+)$; the equality holds in $H(k[\zeta][[X^{\frac{1}{n}}]]^+)$. Formally one can write the above formula as $F_n \cdot \gamma = \sum_{i=0}^{n-1} \gamma(\zeta^i X^{\frac{1}{n}})$.

(2.4) Proposition The following identities hold in $\text{Cart}(k)$.

- (1) $V_1 = F_1 = 1, F_n V_n = n$.
- (2) $[a][b] = [ab]$ for all $a, b \in k$
- (3) $[c]V_n = V_n[c^n], F_n[c] = [c^n]F_n$ for all $c \in k$, all $n \geq 1$.
- (4) $V_m V_n = V_n V_m = V_{mn}, F_m F_n = F_n F_m = F_{mn}$ for all $m, n \geq 1$.
- (5) $F_n V_m = V_m F_n$ if $(m, n) = 1$.
- (6) $(V_n[a]F_n) \cdot (V_m[b]F_m) = r V_{\frac{mn}{r}} [a^{\frac{m}{r}} b^{\frac{n}{r}}] F_{\frac{mn}{r}}, r = (m, n)$, for all $a, b \in k, m, n \geq 1$.

PROOF. The verification of statements (1)–(5) are straightforward using 2.2.1 (i) and left to the reader. The statement (6) follows from (1)–(5). ■

(2.4.1) Exercise Let k be a commutative ring with 1 and let \widetilde{W}^\wedge be the formal completion of the universal Witt vectors, so that $\widetilde{W}^\wedge(N) = 1 + N[[T]] \subset (k \oplus N)[[T]]^\times$.

- (i) Prove that the map which sends every element $\Phi \in \text{End}_k(\widetilde{W}^\wedge)$ to $\Phi_{k[[X]]}(1 - XT)$ establishes a bijection between $\text{End}_k(\widetilde{W}^\wedge)$ with the set of all power series in $k[[X, T]]$ of the form $1 + \sum_{m,n \geq 1} b_{mn} X^m T^n, b_{mn} \in k$.

(ii) Show that $\text{End}_k(\widetilde{W}^\wedge)^{\text{op}}$ can be identified with the set of all expressions

$$\sum_{m,n \geq 1} V_m [a_{mn}] F_n, \quad a_{mn} \in k,$$

such that the endomorphism represented by such a sum sends the element $1 - XT \in \widetilde{W}^\wedge(k[[X]]^+)$ to $\prod_{m,n \geq 1} (1 - a_{mn} X^m T^n)$. All identities in Prop. 2.4 hold in the ring $\text{End}_k(\widetilde{W}^\wedge)^{\text{op}}$.

(2.4.2) Exercise Let k be a commutative ring with 1. The Cartier ring $\text{Cart}(k)$ operates naturally on the *right* of the formal group functor Λ_k . For elements $a, c \in k$ and integers $m, n \geq 1$, prove that

- (i) $(1 - a t^m) \cdot V_n = (1 - a^{\frac{n}{r}} t^{\frac{m}{r}})^r$, where $r = (m, n)$.
- (ii) $(1 - a t^m) \cdot F_n = (1 - a t^{mn})$.
- (iii) $(1 - a t^m) \cdot [c] = (1 - a c^m t^m)$.
- (iv) Deduce 2.4 (6) from (i)–(iii).

(2.5) Proposition *Let k be a commutative ring with 1.)*

(i) *The subset S of $\text{Cart}(k)$ consisting of all elements of the form*

$$\sum_{n \geq 1} V_n [a_n] F_n, \quad a_n \in k \quad \forall n \geq 1$$

form a subring of $\text{Cart}(k)$.

(ii) *The injective map*

$$\widetilde{W}(k) \hookrightarrow \text{Cart}(k), \quad \omega(\underline{a}) \mapsto \sum_{n \geq 1} V_n [a_n] F_n$$

is a homomorphism of rings.

PROOF. Let S' the subset of the power series ring $k[[X, t]]$ consisting of all elements of the form $1 + \sum_{m \geq 1} a_m X^m t^m$ such that $a_m \in k$ for all $m \geq 1$. Clearly S' is a subgroup of the unit group $k[[X, t]]^\times$ of $k[[X, t]]$. By definition $\text{Cart}(k) = \Lambda(k[[X]]^+)$ is a subgroup of $k[[X, t]]^\times$, and $S = S' \cap \text{Cart}(k)$. It follows that S is a subgroup of the additive group underlying $\text{Cart}(k)$. The formula 2.4 (6) implies that the subset $S \subset \text{Cart}(k)$ is stable under multiplication, hence it is a subring. The definition of multiplication for the universal Witt vectors in A.1.1 tells us that the bijection in (ii) is an isomorphism of rings. ■

Corollary Let $A_n(U, V) \in k[U, V]$ be polynomials defined by

$$(1 - UT) \cdot (1 - VT) = (1 - (U + V)T) \cdot \prod_{n \geq 1} (1 - A_n(U, V)T^n)$$

Then for all $c_1, c_2 \in k$ we have

$$[c_1] + [c_2] = [c_1 + c_2] + \sum_{\geq 1} V_n[A_n(c_1, c_2)]F_n.$$

(2.6) Definition The ring $\text{Cart}(k)$ has a natural filtration $\text{Fil}^\bullet \text{Cart}(k)$ by right ideals

$$\text{Fil}^j \text{Cart}(k) = \left\{ \sum_{m \geq j} \sum_{n \geq 1} V_m[a_{mn}]F_n \mid a_{mn} \in k, \forall m \geq j, \exists C_m > 0 \text{ s.t. } a_{mn} = 0 \text{ if } n \geq C_m \right\}$$

for $j \geq 1$. The Cartier ring $\text{Cart}(k)$ is complete with respect to the topology given by the above filtration. Moreover each right ideal $\text{Fil}^j \text{Cart}(k)$ is open and closed in $\text{Cart}(k)$.

(2.7) EXERCISES.

(2.7.1) Exercise Prove the following statements.

- (i) $[c] \cdot \text{Fil}^j \text{Cart}(k) \subseteq \text{Fil}^j \text{Cart}(k)$ for all $c \in k$, all $j \geq 1$.
- (ii) $V_m \cdot \text{Fil}^j \text{Cart}(k) \subseteq \text{Fil}^{mj} \text{Cart}(k)$ for all $m, j \geq 1$.
- (iii) $F_n \cdot \text{Fil}^j \text{Cart}(k) \subseteq \text{Fil}^{\lceil \frac{j}{n} \rceil} \text{Cart}(k)$ for all $n, j \geq 1$.
- (iv) The right ideal of $\text{Cart}(k)$ generated by the elements V_n , $n \geq j$ is dense in $\text{Fil}^j \text{Cart}(k)$.
- (v) The quotient $\text{Cart}(k)/\text{Fil}^2 \text{Cart}(k)$ is canonically isomorphic to k .
- (vi) Left multiplication by V_j induces a bijection

$$V_j : \text{Cart}(k)/\text{Fil}^2 \text{Cart}(k) \xrightarrow{\sim} \text{Fil}^j \text{Cart}(k)/\text{Fil}^{j+1} \text{Cart}(k).$$

(2.7.2) Exercise (i) Show that $\text{Cart}(k)$ is a topological ring, i.e. the multiplication is a continuous map for the topology given by the decreasing filtration $\text{Fil}^\bullet \text{Cart}(k)$ on $\text{Cart}(k)$. (Hint: The point is to show that for any $x \in \text{Cart}(k)$, the map $y \mapsto x \cdot y$ is continuous.)

- (ii) Show that for any $n \geq 1$, there exists $x \in \text{Cart}(k)$ and $y \in \text{Fil}^n \text{Cart}(k)$ such that $x \cdot y \notin \text{Fil}^2 \text{Cart}(k)$.

(2.7.3) Exercise Let k be a commutative ring with 1.

- (i) Show that the right $\text{Cart}(k)$ -module $T := \text{Cart}(k)/\text{Fil}^2\text{Cart}(k)$ is a free k module with basis $x_i :=$ the image of F_i in T .
- (ii) Show that the right $\text{Cart}(k)$ -module T is naturally isomorphic to the tangent space t_Λ of the smooth formal group Λ over k .
- (iii) The right $\text{Cart}(k)$ -module T in (i) above gives a ring homomorphism $\rho : \text{Cart}(k) \rightarrow M'_\infty(k)$, where $M'_\infty(k)$ denotes the set of all $\mathbb{N}_{\geq 1} \times \mathbb{N}_{\geq 1}$ -matrices $(c_{ij})_{i,j \geq 1}$ such that each row has at most a finite number of nonzero entries. Prove that for each element $\sum_{m,n \geq 1} V_m[a_{mn}]F_n \in \text{Cart}(k)$, $\rho(\sum_{m,n \geq 1} V_m[a_{mn}]F_n \in \text{Cart}(k))$ is the matrix $(c_{ij})_{i,j \geq 1}$ with

$$c_{ij} = \sum_{r|(i,j)} \frac{i}{r} \left(a_{\frac{i}{r}, \frac{j}{r}} \right)^r \quad \forall i, j \geq 1.$$
- (iv) Prove that ρ is an injection if and only if the natural map $k \rightarrow k \otimes_{\mathbb{Z}} \mathbb{Q}$ is an injection, or equivalently k is p -torsion free for every prime number p .
- (v) Prove that ρ is an isomorphism if and only if k is a \mathbb{Q} -algebra, or equivalently every nonzero integer is invertible in k .
- (vi) Use (iii) and the properties of the ghost coordinates of the universal Witt vectors to give another proof of 2.5 (ii).

§3. The main theorem of Cartier theory

(3.1) Definition Let k be a commutative ring with 1. A V -reduced $\text{Cart}(k)$ -module is a left $\text{Cart}(k)$ -module M together with a separated decreasing filtration of M

$$M = \text{Fil}^1 M \supset \text{Fil}^2 M \supset \cdots \text{Fil}^n M \supset \text{Fil}^{n+1} M \supset \cdots$$

such that each $\text{Fil}^n M$ is an abelian subgroup of M and

- (i) $(M, \text{Fil}^\bullet M)$ is complete with respect to the topology given by the filtration $\text{Fil}^\bullet M$. In other words, the natural map $\text{Fil}^n M \rightarrow \varprojlim_{m \geq n} (\text{Fil}^n M / \text{Fil}^m M)$ is a bijection for all $n \geq 1$.
- (ii) $V_m \cdot \text{Fil}^n M \subset \text{Fil}^{mn} M$ for all $m, n \geq 1$.
- (iii) The map V_n induces a bijection $V_n : M / \text{Fil}^2 M \xrightarrow{\sim} \text{Fil}^n M / \text{Fil}^{n+1} M$ for every $n \geq 1$.
- (iv) $[c] \cdot \text{Fil}^n M \subset \text{Fil}^n M$ for all $c \in k$ and all $n \geq 1$.
- (v) For every $m, n \geq 1$, there exists an $r \geq 1$ such that $F_m \cdot \text{Fil}^r M \subset \text{Fil}^n M$.

(3.1.1) Definition A V -reduced $\text{Cart}(k)$ -module $(M, \text{Fil}^\bullet M)$ is V -flat if $M / \text{Fil}^2 M$ is a flat k -module. The k -module $M / \text{Fil}^2 M$ is called the tangent space of $(M, \text{Fil}^\bullet M)$.

(3.1.2) As an example, the free $\text{Cart}(k)$ -module $\text{Cart}(k)$ has a filtration with

$$\text{Fil}^n \text{Cart}(k) = \sum_{m \geq n} V_m \text{Cart}(k),$$

making it a V -flat V -reduced $\text{Cart}(k)$ -module. Its tangent space is naturally isomorphic to $k[t]$. See 2.7.1.

(3.1.3) Exercise Let $(M, \text{Fil}^\bullet M)$ be a V -reduced $\text{Cart}(k)$ -module and let n be a positive integer.

- (i) For each $n \geq 1$, the subgroup of M generated by all $V_m \cdot M$, $m \geq n$ is dense in $\text{Fil}^n M$. This follows from 3.1 (i)–(iii).
- (ii) If M is a finitely generated left $\text{Cart}(k)$ -module, then $\text{Fil}^n M = \text{Fil}^n \text{Cart}(k) \cdot M$.
- (iii) Prove that M is finitely generated as a left $\text{Cart}(k)$ -module if and only if $M/\text{Fil}^2 M$ is a finitely generated k -module.
- (iv) Use 2.7.1 to show that properties (iv), (v) in Def. 3.1 follow from 3.1 (i)–(iii).
- (v) Prove the following strengthened form of 3.1 (v):

$$F_m \cdot \text{Fil}^n M \subseteq \text{Fil}^{\lceil \frac{m}{n} \rceil} M \quad \forall m, n \geq 1.$$

(3.1.4) Definition Let $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be a weakly symmetric formal group functor as in 2.1. The abelian group $M_H := H(k[[X]]^+)$ has a natural structure as a left $\text{Cart}(k)$ -module according to Thm. 2.2 The $\text{Cart}(k)$ -module M_H has a natural filtration, with $\text{Fil}^n := \text{Ker}(H(k[[X]]^+ \rightarrow H(k[[X]]^+/X^n k[[X]]))$. We call the pair $(M_H, \text{Fil}^\bullet M_H)$ the Cartier module attached to H .

(3.1.5) Lemma Let $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be weakly symmetric and right exact in the sense of 1.2. Then $(M_H, \text{Fil}^\bullet M_H)$ is a V -reduced $\text{Cart}(k)$ -module.

PROOF. The condition (i), (ii), (iv) of Definition 3.1 are easy to check; it is also easy to verify condition (v) of 3.1 holds with $r = mn$. These are left to the reader as exercises. Here we check that V_n induces an isomorphism from $\text{gr}^1 M_H$ to $\text{gr}^n M_H$ for every $n \geq 1$. Since the functor H is right exact, we have $M_H/\text{Fil}^{n+1} M_H \xrightarrow{\sim} H(k[[X]]^+/X^{n+1} k[[X]])$, and $\text{Fil}^n M_H$ is equal to the image of $H(X^n k[[X]])$ in $H(k[[X]]^+)$ under the map induced by the inclusion $X^n k[[X]] \hookrightarrow k[[X]]^+$. So we have a functorial isomorphism $\text{Fil}^n M_H/\text{Fil}^{n+1} M_H \xrightarrow{\sim} H(X^n k[[X]]/X^{n+1} k[[X]])$ for each $n \geq 1$. The isomorphism

$$k[[X]]^+/X^2 k[[X]] \xrightarrow{\sim} X^n k[[X]]/X^{n+1} k[[X]]$$

in \mathfrak{Nilp}_k which sends X to X^n induces an isomorphism $\text{gr}^1 M_H \xrightarrow{\sim} \text{gr}^n M_H$; this isomorphism is equal to the map induced by V_n . ■

(3.1.6) Lemma Let $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be a group-valued functor. If H is right exact and weakly symmetric, then $(M_H, \text{Fil}^\bullet M_H)$ is a V -reduced $\text{Cart}(k)$ -module. In particular, this is the case if H is a smooth commutative formal group.

PROOF. Exercise. ■

(3.1.7) Definition Let $H : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be weakly symmetric and right exact. We call $(M_H, \text{Fil}^\bullet M_H)$ the V -reduced $\text{Cart}(k)$ -module attached to H . If H is also left exact and commutes with arbitrary direct sums, i.e. H is a smooth commutative formal group, then $(M_H, \text{Fil}^\bullet M_H)$ is V -flat.

(3.1.8) Exercise Let k be a commutative ring with 1. Let $\widetilde{W}(k)$ be the group of universal Witt vectors with entries in k , endowed with the filtration defined in A.1.3 and the action of $\text{Cart}(k)$ defined in A.3. Prove that $(\widetilde{W}(k), \text{Fil}^\bullet \widetilde{W}(k))$ is a V -flat V -reduced $\text{Cart}(k)$ -module. (In fact it is the Cartier module attached to $\widehat{\mathbb{G}}_m$.)

(3.1.9) Exercise Let k be a commutative ring with 1. Let $M = k[[X]]^+$, filtered by $\text{Fil}^n M = X^n k[[X]]$, $n \geq 1$. Define operators $F_n, V_n, [c]$ on M , $n \in \mathbb{N}_{\geq 1}$, $c \in k$ as follows:

- $V_n(\sum_{m \geq 1} a_m X^m) = \sum_{m \geq 1} a_m X^{mn}$,
- $F_n(\sum_{m \geq 1} a_m X^m) = \sum_{m \geq 1} n a_{mn} X^m$,
- $[c](\sum_{m \geq 1} a_m X^m) = \sum_{m \geq 1} a_m c^m X^m$.

Prove that M is a V -reduced V -flat $\text{Cart}(k)$ -module. (In fact it is the Cartier module attached to $\widehat{\mathbb{G}}_a$.)

(3.1.10) Lemma Let $k \rightarrow R$ be a homomorphism between commutative rings with 1. Let $(M, \text{Fil}^\bullet M)$ be a V -reduced $\text{Cart}(k)$ -module. Let $(M_R, \text{Fil}^\bullet M_R)$ be the completion of the pair $(\text{Cart}(R) \otimes_{\text{Cart}(k)} M, \text{Cart}(R) \otimes_{\text{Cart}(k)} \text{Fil}^\bullet M)$.

(i) The pair $(M_R, \text{Fil}^\bullet M_R)$ is a V -reduced $\text{Cart}(R)$ -module.

(ii) If $(M, \text{Fil}^\bullet M)$ is V -flat, then $(M_R, \text{Fil}^\bullet M_R)$ is V -flat.

PROOF. Exercise. ■

(3.1.11) Exercise Let $(M, \text{Fil}^\bullet M)$ be a V -reduced $\text{Cart}(k)$ -module. Let $R = k[\epsilon]/(\epsilon^2)$. The projection $R \rightarrow k$ defines a surjective ring homomorphism $\text{Cart}(R) \rightarrow \text{Cart}(k)$, so we can regard M as a left module over $\text{Cart}(R)$. Show that $(M, \text{Fil}^\bullet M)$ is a V -reduced $\text{Cart}(k)$ -module but it is not a V -flat $\text{Cart}(k)$ -module.

(3.2) Definition Let M be a V -reduced $\text{Cart}(k)$ -module and let Q be a right $\text{Cart}(k)$ -module.

- (i) For every integer $m \geq 1$, let $Q_m := \text{Ann}_Q(\text{Fil}^n \text{Cart}(k))$ be the subgroup of Q consisting of all elements $x \in Q$ such that $x \cdot \text{Fil}^n \text{Cart}(k) = (0)$ for all $n \geq m$. Clearly we have $Q_1 \subseteq Q_2 \subseteq Q_3 \subseteq \dots$.
- (ii) For each $m, r \geq 1$, define $Q_m \odot M^r$ to be the image of $Q_m \otimes \text{Fil}^r M$ in $Q \otimes_{\text{Cart}(k)} M$. If $r \geq m$ and $s \geq m$, then $Q_m \odot M^r = Q_m \odot M^s$; see 3.2.1. Hence $Q_m \odot M^m \subseteq Q_n \odot M^n$ if $m \leq n$.
- (iii) Define the reduced tensor product $Q \overline{\otimes}_{\text{Cart}(k)} M$ by

$$Q \overline{\otimes}_{\text{Cart}(k)} M = Q \otimes_{\text{Cart}(k)} M \Big/ \left(\bigcup_m (Q_m \odot M^m) \right).$$

(3.2.1) Exercise Notation as in 3.2.

- (i) For every $x \in Q_m$ and every $y \in \text{Fil}^n M$ with $n \geq m$, let $y_1 \in M$ and $y_2 \in \text{Fil}^{n+1} M$ be such that $y = V^n y_1 + y_2$. Then $x \otimes y = x \otimes y_2$ in $Q \otimes_{\text{Cart}(k)} M$.
- (ii) Show that $Q_m \odot M^r = Q_m \odot M^s$ if $r, s \geq m$.

(3.2.2) Exercise Let $(M, \text{Fil}^\bullet M)$ be a V -reduced $\text{Cart}(k)$ -module.

- (i) Let N be a nilpotent k -algebra such that $N^2 = (0)$. Prove that $\Lambda(N) \overline{\otimes}_{\text{Cart}(k)} M \cong N \otimes_k (M/\text{Fil}^2 M)$.
- (ii) Prove that $\Lambda(k[[X]]^+/X^n k[[X]]) \overline{\otimes}_{\text{Cart}(k)} M \cong M/\text{Fil}^n M$.

(3.3) Theorem *Let k be a commutative ring with 1. Then there is a canonical equivalence of categories, between the category of smooth commutative formal groups over k as defined in 1.3 and the category of V -flat V -reduced $\text{Cart}(k)$ -modules, defined as follows.*

$$\begin{array}{ccc} \{\text{smooth formal groups over } k\} & \xrightarrow{\sim} & \{V\text{-flat } V\text{-reduced } \text{Cart}(k)\text{-mod}\} \\ G & \xrightarrow{\quad} & M_G = \text{Hom}(\Lambda, G) \\ \Lambda \overline{\otimes}_{\text{Cart}(k)} M & \xleftarrow{\quad} & M \end{array}$$

Recall that $M_G = \text{Hom}(\Lambda, G)$ is canonically isomorphic to $G(X k[[X]])$, the group of all formal curves in the smooth formal group G . The reduced tensor product $\Lambda \overline{\otimes}_{\text{Cart}(k)} M$ is the functor whose value at any nilpotent k -algebra N is $\Lambda(N) \overline{\otimes}_{\text{Cart}(k)} M$.

The key steps of the proof of Thm. 3.3 are Prop. 3.3.3 and Thm. 3.3.5.

(3.3.1) Lemma *Let $\alpha : (L, \text{Fil}^\bullet L) \rightarrow (M, \text{Fil}^\bullet M)$ be a homomorphism between V -reduced $\text{Cart}(k)$ -modules, i.e. $\alpha(\text{Fil}^i L) \subseteq \text{Fil}^i M$ for all $i \geq 1$. Then the following are equivalent.*

(i) $\alpha(\text{Fil}^i L) = \text{Fil}^i M$ for all $i \geq 1$.

(ii) $\alpha(L) = M$.

(iii) $\alpha : L/\text{Fil}^2 L \rightarrow M/\text{Fil}^2 M$ is surjective.

(3.3.2) Exercise Let k be a commutative ring with 1. Let I be any set. Denote $\text{Cart}(k)^{(I)}$ the free $\text{Cart}(k)$ -module with basis I , and give it the filtration such that $\text{Fil}^i \text{Cart}(k)^{(I)} = (\text{Fil}^i \text{Cart}(k))^{(I)}$.

(i) Show that $(\text{Cart}(k)^{(I)}, \text{Fil}^\bullet \text{Cart}(k)^{(I)})$ is a V -reduced $\text{Cart}(k)$ -module if and only if I is finite.

(ii) Let $\widehat{\text{Cart}(k)^{(I)}}$ be the completion of the filtered module. $(\text{Cart}(k)^{(I)}, \text{Fil}^\bullet \text{Cart}(k)^{(I)})$, with the induced filtration. Prove that $\widehat{\text{Cart}(k)^{(I)}}$ is a V -reduced $\text{Cart}(k)$ -module. We call $\widehat{\text{Cart}(k)^{(I)}}$ the free V -reduced $\text{Cart}(k)$ -module with basis I . Formulate a universal property which justifies this terminology.

(iii) Let Q be a torsion right $\text{Cart}(k)$ -module, i.e. $Q = \bigcup_n \text{Ann}_Q(\text{Fil}^n \text{Cart}(k))$. Prove that $Q \widehat{\otimes}_{\text{Cart}(k)} \widehat{\text{Cart}(k)^{(I)}}$ is naturally isomorphic to $Q^{(I)}$.

(3.3.3) Proposition Let $\alpha : (L, \text{Fil}^\bullet L) \rightarrow (M, \text{Fil}^\bullet M)$ be a surjective homomorphism between V -reduced $\text{Cart}(k)$ -modules as in Lemma 3.3.1. Let K be the kernel of α , with the induced filtration $\text{Fil}^i K = K \cap \text{Fil}^i L$ for all $i \geq 1$. Then $(K, \text{Fil}^\bullet K)$ is a V -reduced $\text{Cart}(k)$ -module.

PROOF. We have a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K/\text{Fil}^2 K & \longrightarrow & L/\text{Fil}^2 L & \xrightarrow{\alpha} & M/\text{Fil}^2 M & \longrightarrow & 0 \\ & & \downarrow V_n & & \cong \downarrow V_n & & \cong \downarrow V_n & & \\ 0 & \longrightarrow & \text{Fil}^n K/\text{Fil}^{n+1} K & \longrightarrow & \text{Fil}^n L/\text{Fil}^{n+1} L & \xrightarrow{\alpha} & \text{Fil}^n M/\text{Fil}^{n+1} M & \longrightarrow & 0 \end{array}$$

with exact rows. From the five-lemma we see that V_n induces a bijection $V_n : K/\text{Fil}^2 K \xrightarrow{\sim} \text{Fil}^n K/\text{Fil}^{n+1} K$ for all $n \geq 1$. The rest of the conditions for $(K, \text{Fil}^\bullet K)$ to be V -reduced is easy. ■

(3.3.4) Definition Prop. 3.3.3 shows that for every V -reduced $\text{Cart}(k)$ -module $(M, \text{Fil}^\bullet M)$, there exists a free resolution

$$\cdots \rightarrow L_i \xrightarrow{\partial_i} L_{i-1} \xrightarrow{\partial_{i-1}} \cdots \xrightarrow{\partial_2} L_1 \xrightarrow{\partial_1} L_0 \xrightarrow{\partial_0} M \rightarrow 0$$

of M , where each L_i is a free V -reduced $\text{Cart}(k)$ -modules in the sense of 3.3.2 (ii), each ∂_i is compatible with the filtrations, and $\text{Ker}(\partial_i) = \text{Image}(\partial_{i+1})$ for all $i \geq 0$, and ∂_0 is surjective. Define *reduced torsion functors* $\overline{\text{Tor}}_i^{\text{Cart}(k)}(?, M)$ by

$$\overline{\text{Tor}}_i^{\text{Cart}(k)}(Q, M) = H_i \left(Q \overline{\otimes}_{\text{Cart}(k)} (\cdots \rightarrow L_i \xrightarrow{\partial_i} L_{i-1} \xrightarrow{\partial_{i-1}} \cdots \xrightarrow{\partial_2} L_1 \xrightarrow{\partial_1} L_0) \right)$$

for any torsion right $\text{Cart}(k)$ -module Q .

Exercise (i) Prove that the functor $\overline{\text{Tor}}_i^{\text{Cart}(k)}$ is well-defined.

(ii) Show that every short exact sequence of torsion right $\text{Cart}(k)$ -modules gives rise to a long exact sequence for the functor $\overline{\text{Tor}}^{\text{Cart}(k)}$.

(iii) Formulate and prove a similar statement for the second entry of the reduced torsion functor.

(3.3.5) Theorem *let k be a commutative ring with 1. Let $(M, \text{Fil}^\bullet M)$ be a V -reduced $\text{Cart}(k)$ -module. Let N be a nilpotent k -algebra N .*

(i) *Suppose that $(M, \text{Fil}^\bullet M)$ is V -flat, i.e. $M/\text{Fil}^2 M$ is a flat k -module. Then*

$$\overline{\text{Tor}}_i^{\text{Cart}(k)}(\Lambda(N), M) = (0)$$

for all $i \geq 1$.

(ii) *Suppose that N has a finite decreasing filtration*

$$N = \text{Fil}^1 N \supseteq \text{Fil}^2 N \supseteq \cdots \supseteq \text{Fil}^s N = (0)$$

such that each $\text{Fil}^j N$ is an ideal of N , $(\text{Fil}^j N)^2 \subseteq \text{Fil}^{j+1} N$ and $\text{Fil}^j N/\text{Fil}^{j+1} N$ is a flat k -module for $j = 1, \dots, s-1$. Then $\overline{\text{Tor}}_i^{\text{Cart}(k)}(\Lambda(N), M) = 0$ for all $i \geq 1$.

PROOF OF (I). Choose an $s \in \mathbb{N}$ such that $N^s = 0$. Then we have a decreasing filtration

$$\Lambda(N) \supseteq \Lambda(N^2) \supseteq \cdots \supseteq \Lambda(N^{s-1}) \supseteq \Lambda(N^s) = (0)$$

of $\Lambda(N)$. For each $j = 1, 2, \dots, s-1$ we have $\Lambda(N^j)/\Lambda(N^{j+1}) \cong \Lambda(N^j/N^{j+1})$ as right $\text{Cart}(k)$ -modules, hence it suffices to show that $\overline{\text{Tor}}_i^{\text{Cart}(k)}(\Lambda(N^j/N^{j+1}), M) = 0$ for each $j = 1, 2, \dots, s-1$. Let $L \twoheadrightarrow M$ be a surjection from a free reduced $\text{Cart}(k)$ -module L to M , and let K be the kernel. Then K is also a V -flat V -reduced $\text{Cart}(k)$ -module. Recall from Exer. 3.2.2 (i) that $\Lambda(N^j/N^{j+1}) \overline{\otimes}_{\text{Cart}(k)} M \cong (N^j/N^{j+1}) \otimes_k (M/\text{Fil}^2 M)$ for each $j = 1$. The long exact sequence attached to the short exact sequence $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$ yields isomorphisms

$$\overline{\text{Tor}}_{i+1}^{\text{Cart}(k)}(\Lambda(N^j/N^{j+1}), M) \xrightarrow{\sim} \overline{\text{Tor}}_i^{\text{Cart}(k)}(\Lambda(N^j/N^{j+1}), K), \quad i \geq 1$$

and an exact sequence

$$(N^j/N^{j+1}) \otimes_k (K/\text{Fil}^2 K) \xrightarrow{\alpha} (N^j/N^{j+1}) \otimes_k (L/\text{Fil}^2 L) \rightarrow (N^j/N^{j+1}) \otimes_k (M/\text{Fil}^2 M) \rightarrow 0$$

such that the kernel of α is isomorphic to $\overline{\text{Tor}}_1^{\text{Cart}(k)}(\Lambda(N^j/N^{j+1}), M)$. Since $M/\text{Fil}^2 M$ is a flat k -module, we see that $\overline{\text{Tor}}_1^{\text{Cart}(k)}(\Lambda(N^j/N^{j+1}), M) = 0$ for every V -flat V -reduced $\text{Cart}(k)$ -module M . Since K is also V -flat, $\overline{\text{Tor}}_2^{\text{Cart}(k)}(\Lambda(N^j/N^{j+1}), M) = 0$ as well. An induction shows that $\overline{\text{Tor}}_i^{\text{Cart}(k)}(\Lambda(N^j/N^{j+1}), M) = 0$ for all $i \geq 1$ and all $j = 1, 2, \dots, s-1$. The statement (i) follows.

PROOF OF (II). In the proof of (i) above, replace the ideals N^j by $\text{Fil}^j N$. The sequence

$$\begin{aligned} (\text{Fil}^j N/\text{Fil}^{j+1} N) \otimes_k (K/\text{Fil}^2 K) &\xrightarrow{\alpha} (\text{Fil}^j N/\text{Fil}^{j+1} N) \otimes_k (L/\text{Fil}^2 L) \\ &\rightarrow (\text{Fil}^j N/\text{Fil}^{j+1} N) \otimes_k (M/\text{Fil}^2 M) \rightarrow 0 \end{aligned}$$

is exact because $\text{Fil}^j N/\text{Fil}^{j+1} N$ is a flat k -module. The rest of the proof of (ii) is the same as the proof of (i). ■

PROOF OF THM. 3.3. Suppose that $(M, \text{Fil}^\bullet M)$ is a V -flat V -reduced $\text{Cart}(k)$ -module. It follows immediately from Thm. 3.3.5 that $G := \Lambda \overline{\otimes}_{\text{Cart}(k)} M$ is a smooth formal group. Conversely given any smooth formal group G , M_G is V -reduced according to Lemma 3.1.6. By Exer. 3.2.2 (ii), we have a functorial isomorphism

$$(\Lambda \overline{\otimes}_{\text{Cart}(k)} M)(k[[X]]^+) = \text{proj} \lim_n \Lambda(k[[X]]^+/X^n k[[X]]) \overline{\otimes}_{\text{Cart}(k)} M \xrightarrow{\sim} M$$

for each V -flat V -reduced $\text{Cart}(k)$ -module M .

To finish the proof, it remains to produce a functorial isomorphism

$$\beta_G : \Lambda \overline{\otimes}_{\text{Cart}(k)} M_G \xrightarrow{\sim} G$$

for each smooth formal group G . For each nilpotent k -algebra N , we have a natural map $\beta_{G,N} : \Lambda \overline{\otimes}_{\text{Cart}(k)} M_G \rightarrow G$ such that $\beta_{G,N} : \sum_i f_i \otimes h_i \mapsto \sum_i \Phi_{h_i}(f_i) \in G(N)$ in the notation of Cor. 2.2.1, where $f_i \in \Lambda(N)$ and $h_i \in M_G$ for each i . The map $\beta_{G,N}$ factors through $\Lambda(N) \overline{\otimes}_{\text{Cart}(k)} M_G$ by 2.2.1 (iv) and gives the desired map $\alpha_{G,N} : \Lambda(N) \overline{\otimes}_{\text{Cart}(k)} M_G \rightarrow G(N)$. Since both the source and the target of β_G are exact and commutes with arbitrary direct sums, to show that β_G is an isomorphism for every nilpotent k -algebra N it suffices to verify this statement when $N^2 = (0)$ and N is isomorphic to k as a k -module. In that case $\beta_{G,N}$ is the canonical isomorphism $M_G/\text{Fil}^2 M_G \xrightarrow{\sim} \mathfrak{t}_G$. ■

(3.3.6) Exercise Prove that the equivalence of categories in Thm. 3.3 extends to an equivalence of categories between the category of V -reduced V -flat $\text{Cart}(k)$ -modules and the category of functors $G : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ which are right exact and weakly left exact.

(3.3.7) Exercise Let M be a V -reduced V -flat $\text{Cart}(k)$ -module. Let k' be a commutative k algebra with 1. Let $M' = \text{Cart}(k') \widehat{\otimes}_{\text{Cart}(k)} M$, defined as the completion of the left $\text{Cart}(k)$ -module $\text{Cart}(k') \otimes_{\text{Cart}(k)} M$ with respect to the filtration given by the image of $\text{Fil}^\bullet \text{Cart}(k') \otimes_{\text{Cart}(k)} M$ in $\text{Cart}(k') \otimes_{\text{Cart}(k)} M$, endowed with the induced filtration.

- (i) The pair $(M', \text{Fil}^\bullet M')$ is V -reduced, and $k' \otimes_k (M/\text{Fil}^2 M) \xrightarrow{\sim} M'/\text{Fil}^2 M'$ as k' -modules.
- (ii) For every nilpotent k' -algebra N , there is a canonical isomorphism

$$\Lambda(N) \overline{\otimes}_{\text{Cart}(k)} M \xrightarrow{\sim} \Lambda(N) \overline{\otimes}_{\text{Cart}(k')} M'.$$

§4. Localized Cartier theory

In this section we fix a prime number p . Let k be a commutative ring with 1 over $\mathbb{Z}_{(p)}$.

(4.1) Definition Recall from 2.3.4 that every prime number $\ell \neq p$ is invertible in $\text{Cart}(k)$. Define elements ϵ_p and $\epsilon_{p,n}$ of the Cartier ring $\text{Cart}(k)$ for $n \in \mathbb{N}$, $(n, p) = 1$ by

$$\begin{aligned} \epsilon_p = \epsilon_{p,1} &= \sum_{\substack{(n,p)=1 \\ n \geq 1}} \frac{\mu(n)}{n} V_n F_n = \prod_{\substack{\ell \neq p \\ \ell \text{ prime}}} \left(1 - \frac{1}{\ell} V_\ell F_\ell \right) \\ \epsilon_{p,n} &= \frac{1}{n} V_n \epsilon_p F_n \end{aligned}$$

In the above μ is the Möbius function on $\mathbb{N}_{\geq 1}$, characterized by the following properties: $\mu(mn) = \mu(m)\mu(n)$ if $(m, n) = 1$, and for every prime number ℓ we have $\mu(\ell) = -1$, $\mu(\ell^i) = 0$ if $i \geq 2$.

(4.1.1) Proposition *The following properties hold.*

- (i) $\epsilon_p^2 = \epsilon_p$.
- (ii) $\sum_{\substack{(n,p)=1 \\ n \geq 1}} \epsilon_{p,n} = 1$.
- (iii) $\epsilon_p V_n = 0$, $F_n \epsilon_p = 0$ for all n with $(n, p) = 1$.
- (iv) $\epsilon_{p,n}^2 = \epsilon_{p,n}$ for all $n > 1$ with $(n, p) = 1$.
- (v) $\epsilon_{p,n} \epsilon_{p,m} = 0$ for all $m \neq n$ with $(mn, p) = 1$.
- (vi) $[c] \epsilon_p = \epsilon_p [c]$ and $[c] \epsilon_{p,n} = \epsilon_{p,n} [c]$ for all $c \in k$ and all n with $(n, p) = 1$.
- (vii) $F_p \epsilon_{p,n} = \epsilon_{p,n} F_p$, $V_p \epsilon_{p,n} = \epsilon_{p,n} V_p$ for all n with $(n, p) = 1$.

PROOF. From 2.4 (1)–(5), one easily deduces that for every prime number $\ell \neq p$ we have

$$\left(1 - \frac{1}{\ell}V_\ell F_\ell\right)V_\ell = 0, \quad F_\ell\left(1 - \frac{1}{\ell}V_\ell F_\ell\right) = 0, \quad \text{and} \left(1 - \frac{1}{\ell}V_\ell F_\ell\right)^2 = \left(1 - \frac{1}{\ell}V_\ell F_\ell\right).$$

The statements (i) and (iii) follows. Statement (v) is an easy consequence of (iii). The proof of statement (iv) is an easy computation:

$$\left(\frac{1}{n}V_n\epsilon_p F_n\right)^2 = \frac{1}{n^2}V_n\epsilon_p F_n V_n\epsilon_p F_n = \frac{1}{n}V_n\epsilon_p F_n.$$

By 2.4 (4) and (5), the statement (ii) is a consequence of the following telescoping identity:

$$\sum_{m \geq 0} \frac{1}{\ell^m} V_{\ell^m} \left(1 - \frac{1}{\ell}V_\ell F_\ell\right) F_{\ell^m} = 1$$

for any prime number $\ell \neq p$.

To prove (vi), observe first that $[c]\epsilon_p = \epsilon_p[c]$ by 2.4 (3); this in turn gives

$$[c]\epsilon_{p,n} = \frac{1}{n}V_n[c^n]\epsilon_p F_n = \frac{1}{n}V_n\epsilon_p[c^n]F_n = \epsilon_p[c].$$

■

(4.1.2) Definition (i) Denote by $\text{Cart}_p(k)$ the subring $\epsilon_p \text{Cart}(k) \epsilon_p$ of $\text{Cart}(k)$. Note that ϵ_p is the unit element of $\text{Cart}_p(k)$.

(ii) Define elements $F, V \in \text{Cart}_p(k)$ by

$$F = \epsilon_p F_p = F_p \epsilon_p = \epsilon_p F_p \epsilon_p, \quad V = \epsilon_p V_p = V_p \epsilon_p = \epsilon_p V_p \epsilon_p.$$

(iii) For every element $c \in k$, denote by $\langle c \rangle$ the element $\epsilon_p[c]\epsilon_p = \epsilon_p[c] = [c]\epsilon_p \in \text{Cart}_p(k)$.

(4.1.3) Exercise Let $E(T) \in \mathbb{Q}[[T]]$ be the power series

$$E(T) = \prod_{\substack{n \in \mathbb{N} \\ (n,p)=1}} (1 - T^n)^{\frac{\mu(n)}{n}} = \exp\left(-\sum_{m \geq 0} \frac{T^{p^m}}{p^m}\right).$$

(i) Verify the second equality in the displayed formula above for $E(T)$, and prove that all coefficients of $E(T)$ lie in $\mathbb{Z}_{(p)}$.

(ii) Recall that the additive group underlying $\text{Cart}(k)$ is a subgroup of $k[[X, t]]^\times$ by definition. Show that for any element $x = \sum_{m,n \geq 1} V^m[a_{mn}]F^n$ in $\text{Cart}(k)$ with $a_{mn} \in k$ for all $m, n \geq 1$, the element $\epsilon_p x \epsilon_p$ is represented by the element

$$\prod_{m,n \geq 1} E(a_{mn} X^{p^m} t^{p^n}).$$

(4.1.4) Exercise Notation as above. Prove that for any left $\text{Cart}(k)$ -module M , the subgroup $\epsilon_p(M)$ consists of all elements $x \in M$ such that $F_n x = 0$ for all $n > 1$ with $(n, p) = 1$. Elements of M with the above property will be called *p-typical elements*.

(4.1.5) Exercise Prove the following identities in $\text{Cart}_p(k)$.

- (1) $F \langle a \rangle = \langle a^p \rangle F$ for all $a \in k$.
- (2) $\langle a \rangle V = V \langle a^p \rangle$ for all $a \in k$.
- (3) $\langle a \rangle \langle b \rangle = \langle ab \rangle$ for all $a, b \in k$.
- (4) $FV = p$.
- (5) $VF = p$ if and only if $p = 0$ in k .
- (6) Every prime number $\ell \neq p$ is invertible in $\text{Cart}_p(k)$. The prime number p is invertible in $\text{Cart}_p(k)$ if and only if p is invertible in k .
- (7) $V^m \langle a \rangle F^m V^n \langle b \rangle F^n = p^r V^{m+n-r} \langle a^{p^{n-r}} b^{p^{m-r}} \rangle F^{m+n-r}$ for all $a, b \in k$ and all $m, n \in \mathbb{N}$, where $r = \min\{m, n\}$.

(4.2) Definition Let k be a commutative $\mathbb{Z}_{(p)}$ -algebra with 1. Denote by Λ_p the image of Λ_p . In other words, Λ_p is a functor from \mathfrak{Nilp}_k to \mathfrak{Ab} such that

$$\Lambda_p(N) = \Lambda(N) \cdot \epsilon_p$$

for any nilpotent k -algebra N .

(4.2.1) Exercise Let $E(T) \in \mathbb{Z}_{(p)}[[T]]$ be the inverse of the Artin-Hasse exponential as in 4.1.2.

- (i) Prove that for any nilpotent k -algebra N , every element of $\Lambda_p(N)$ has a unique expression as a finite product

$$\prod_{i=0}^m E(u_i t^{p^i})$$

for some $m \in \mathbb{N}$, and $u_i \in N$ for $i = 0, 1, \dots, m$.

- (ii) Prove that Λ_p is a smooth commutative formal group over k .

(4.2.2) Proposition (i) *The local Cartier ring $\text{Cart}_p(k)$ is complete with respect to the decreasing sequence of right ideals $V^i \text{Cart}_p(k)$.*

- (ii) *Every element of $\text{Cart}_p(k)$ can be expressed as a convergent sum*

$$\sum_{m,n \geq 0} V^m \langle a_{mn} \rangle F^n, \quad a_{mn} \in k, \forall m \exists C_m > 0 \text{ s.t. } a_{mn} = 0 \text{ if } n \geq C_m.$$

(iii) The set of all elements of $\text{Cart}_p(k)$ which can be represented as a convergent sum of the form

$$\sum_{m \geq 0} V^m \langle a_m \rangle F^m, \quad a_m \in k$$

is a subring of $\text{Cart}_p(k)$. The map

$$w_p(\underline{a}) \mapsto \sum_{m \geq 0} V^m \langle a_m \rangle F^m \quad \underline{a} = (a_0, a_1, a_2, \dots), \quad a_i \in k \quad \forall i \geq 0$$

establishes an isomorphism from the ring of p -adic Witt vectors $W_p(k)$ to the above subring of $\text{Cart}_p(k)$.

PROOF. Statement (i) and the existence part of statement (ii) are easy and left as an exercises. To prove the uniqueness part of (ii), according to 4.1.3 it suffices to check that if $(a_{mn})_{m,n \in \mathbb{N}}$ is a family of elements in k such that the infinite product

$$\prod_{m,n \geq 1} E(a_{mn} X^{p^m} t^{p^n})$$

is equal to 1 in $k[[X, t]]$, then $a_{mn} = 0$ for all $m, n \geq 1$. This follows from the fact that

$$E(X) \equiv 1 + X \pmod{(X^p)}.$$

The statement (iii) follows from 4.1.5 (7) and the property 4.1.5 of multiplication in the ring of p -adic Witt vectors. ■

(4.2.3) Exercise Let $j_p : \Lambda_p \rightarrow \Lambda$ be the homomorphism of smooth commutative formal groups over k induced by the inclusion map, and let $\pi_p : \Lambda \rightarrow \Lambda_p$ be the homomorphism induced by ϵ_p . Let $G : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ be a functor. The abelian group $\text{Hom}(\Lambda, G)$ has a natural structure as a left module over $\text{Cart}(k) = \text{End}(\Lambda/k)^{\text{op}}$.

- (i) Prove that for every homomorphism $h \in \text{Hom}(\Lambda_p, G)$ $h \circ \pi_p \in \text{Hom}(\Lambda, G)$ is a p -typical element of $\text{Hom}(\Lambda, G)$.
- (ii) Prove that the map $h \mapsto h \circ \pi_p$ above establishes a bijection from $\text{Hom}(\Lambda_p, G)$ to the set of all p -typical elements in $\text{Hom}(\Lambda, G)$, whose inverse is given by $h' \mapsto h' \circ j_p$.

(4.2.4) Exercise Prove that $\text{Cart}_p(k)$ is naturally isomorphic to $\text{End}(\Lambda_p)^{\text{op}}$, the opposite ring of the endomorphism ring of $\text{End}(\Lambda_p)$.

(4.2.5) Exercise Let k be a commutative algebra over $\mathbb{Z}_{(p)}$. Let T be the right $\text{Cart}_p(k)$ -module $\text{Cart}_p(k)/V\text{Cart}_p(k)$.

- (i) Show that there is a natural isomorphism from T the tangent space of the smooth commutative formal group $\Lambda_p : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$.

- (ii) Show that the element $x_i :=$ the image of F^i in T , $i = 0, 1, 2, \dots$, form a k -basis of T .
- (iii) The basis x_i , $i \in \mathbb{N}$ of the right $\text{Cart}_p(k)$ -module T defines a ring homomorphism $\rho_p : \text{Cart}_p(k) \rightarrow M'_{\mathbb{N}}(k)$, where $M'_{\mathbb{N}}(k)$ consists of all $\mathbb{N} \times \mathbb{N}$ -matrices $(c_{ij})_{i,j \geq 0}$ with at most finitely many non-zero entries in each row, and $c_{ij} \in k$ for all $i, j \geq 0$. Let $x = \sum_{m,n \geq 0} V^m \langle a_{mn} \rangle F^n$ be an element of $\text{Cart}_p(k)$, $a_{mn} \in k$ for all $m, n \geq 0$. Prove that $\rho_p(x)$ is the matrix (c_{ij}) whose (i, j) -th entry is given by

$$c_{ij} = \sum_{r \leq \min\{i,j\}} p^{i-r} (a_{i-r, j-r})^{p^r}$$

for all $i, j \geq 0$.

- (iii) Use the formula in (ii) above to give another proof of 4.2.2 (iii).

(4.3) Definition Let k be a commutative $\mathbb{Z}_{(p)}$ -algebra.

- (i) A V -reduced $\text{Cart}_p(k)$ -module M is a left $\text{Cart}_p(k)$ -module such that the map $V : M \rightarrow M$ is injective and the canonical map $M \rightarrow \varprojlim_n (M/V^n M)$ is an isomorphism.
- (ii) A V -reduced $\text{Cart}_p(k)$ -module M is V -flat if M/VM is a flat k -module.

(4.3.1) Definition Let I be a set. Denote by $\text{Cart}_p(k)^{(I)}$ the direct sum of copies of $\text{Cart}_p(k)$ indexed by I . The completion of the free $\text{Cart}_p(k)$ -module $\text{Cart}_p(k)^{(I)}$ with respect to the filtered family of subgroups $V^i \text{Cart}_p(k)^{(I)}$ is a V -reduced $\text{Cart}_p(k)$ -module, denoted by $\widehat{\text{Cart}_p(k)^{(I)}}$; we called it the free V -reduced $\text{Cart}_p(k)$ -module with basis indexed by I .

(4.3.2) Lemma Every element of the subset $\text{Cart}(k)\epsilon_p$ of $\text{Cart}(k)$ can be expressed as a convergent sum

$$\sum_{(n,p)=1} V_n x_n, \quad x_n \in \text{Cart}_p(k) \quad \forall n \text{ with } (n, p) = 1$$

for uniquely determined elements $x_n \in \text{Cart}_p(k)$, $(n, p) = 1$. Conversely every sequence of elements $(x_n)_{(n,p)=1}$ in $\text{Cart}_p(k)$ defines an element of $\text{Cart}(k)\epsilon_p$:

$$\text{Cart}(k)\epsilon_p = \widehat{\bigoplus_{(n,p)=1} V_n \cdot \text{Cart}_p(k)}.$$

PROOF. For $x \in \text{Cart}(k)\epsilon_p$, we have

$$x = \sum_{(n,p)=1} \epsilon_{p,n} \cdot x = \sum_{(n,p)=1} V_n \left(\frac{1}{n} \epsilon_p F_n x \epsilon_p \right).$$

On the other hand, suppose that we have $\sum_{(n,p)=1} V_n x_n = 0$ and $x_n \in \text{Cart}_p(k)$ for all $n \geq 1$ with $(n,p) = 1$. For any $m \geq 1$ with $(m,p) = 1$, we have

$$0 = \epsilon_{p,m} \cdot \sum_{(n,p)=1} V_n x_n = V_m x_m$$

because $\epsilon_p V_r = 0$ and $F_r x_n = 0$ for all $r > 1$ with $(r,p) = 1$ and all $n \geq 1$ with $(n,p) = 1$. Hence $x_m = 0$ since left multiplication by V_m on $\text{Cart}(k)$ is injective. ■

(4.3.3) Lemma *Let M_p be a V -reduced $\text{Cart}_p(k)$ -module. Let $\text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p$ be the V -adic completion of the tensor product $\text{Cart}(k)\epsilon_p \otimes_{\text{Cart}_p(k)} M_p$, defined as the completion of $\text{Cart}(k)\epsilon_p \otimes_{\text{Cart}_p(k)} M_p$ with respect to the decreasing family of the subgroups*

$$\text{Image}(\text{Fil}^i \text{Cart}(k)\epsilon_p \otimes_{\text{Cart}_p(k)} M_p \rightarrow \text{Cart}(k)\epsilon_p \otimes_{\text{Cart}_p(k)} M_p)$$

of $\text{Cart}(k)\epsilon_p \otimes_{\text{Cart}_p(k)} M_p$.

(i) *The completed tensor product $\text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p$ is the topological direct sum*

$$\widehat{\bigoplus}_{(n,p)=1} V_n \otimes M_p$$

of its subgroups $V_n \otimes M_p$, $(n,p) = 1$, so that we have a natural bijection between $\text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p$ and the set of sequences $(x_n)_{(n,p)=1}$ of elements in M_p indexed by positive integers prime to p .

(ii) *Define a decreasing filtration on the completed tensor product by*

$$\text{Fil}^m \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p = \overline{\text{Image}(\text{Fil}^m \text{Cart}(k)\epsilon_p \otimes M_p \rightarrow \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p)},$$

the closure of the image of $\text{Fil}^m \text{Cart}(k)\epsilon_p \otimes M_p$, $m \geq 1$. Then the completed tensor product $\text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p$ is a V -reduced $\text{Cart}(k)$ -module.

(iii) *The inclusion map $M_p \hookrightarrow \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p$ induces an isomorphism*

$$M_p / VM_p \xrightarrow{\sim} \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p / \text{Fil}^2 \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p.$$

(iv) *There is a canonical isomorphism*

$$\epsilon_p \cdot \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p \xrightarrow{\sim} M_p.$$

from the set of all p -typical elements in $\text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p$ to M_p .

PROOF. This Lemma is a corollary of 4.3.2. The isomorphism in (iv) is

$$\epsilon_p \cdot \left(\sum_{(n,p)=1} V_n \otimes x_n \right) \mapsto x_1,$$

whose inverse is induced by the inclusion. ■

(4.3.4) Lemma *Let M be a V -reduced $\text{Cart}(k)$ module and let $M_p = \epsilon_p M$ be the set of all p -typical elements in M .*

(i) *The canonical map*

$$\widehat{\bigoplus_{\substack{(n,p)=1 \\ n \geq 1}} M_p} \longrightarrow M \quad (1)$$

$$(x_n)_{(n,p)=1} \mapsto \sum_{(n,p)=1} V_n x_n \quad (2)$$

is an isomorphism.

(ii) *The canonical map $\text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p \rightarrow M$ is an isomorphism.*

PROOF. The argument of 4.3.2 proves (i). The statement (ii) follows from (i) and 4.3.3.

Combining 4.3.2, 4.3.3 and 4.3.4, we obtain the following theorem.

(4.4) Theorem *Let k be a commutative $\mathbb{Z}_{(p)}$ -algebra with 1.*

(i) *There is an equivalence of categories between the category of V -reduced $\text{Cart}(k)$ -modules and the category of V -reduced $\text{Cart}_p(k)$ -modules, defined as follows.*

$$\begin{array}{ccc} \{ V\text{-reduced } \text{Cart}(k)\text{-mod} \} & \xrightarrow{\sim} & \{ V\text{-reduced } \text{Cart}_p(k)\text{-mod} \} \\ M & \xrightarrow{\quad \quad \quad} & \epsilon_p M \\ \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p & \xleftarrow{\quad \quad \quad} & M_p \end{array}$$

(ii) *Let M be a V -reduced $\text{Cart}(k)$ -module M , and let M_p be the V -reduced $\text{Cart}_p(k)$ -module M_p attached to M as in (i) above. Then there is a canonical isomorphism $M/\text{Fil}^2 M \cong M_p/V M_p$. In particular M is V -flat if and only if M_p is V -flat. Similarly M is a finitely generated $\text{Cart}(k)$ -module if and only if M_p is a finitely generated $\text{Cart}_p(k)$ -module.*

The next theorem is the local version of the main theorem of Cartier theory. The main ingredients of the proof occupies 4.5.1–4.5.6, and the end of the proof is in 4.5.7.

(4.5) Theorem *Let k be a commutative $\mathbb{Z}_{(p)}$ -algebra with 1. Then there is a canonical equivalence of categories, between the category of smooth commutative formal groups over k as defined in 1.3 and the category of V -flat V -reduced $\text{Cart}_p(k)$ -modules, defined as follows.*

$$\begin{array}{ccc} \{\text{smooth formal groups over } k\} & \xrightarrow{\sim} & \{V\text{-flat } V\text{-reduced } \text{Cart}_p(k)\text{-mod}\} \\ G & \xrightarrow{\quad \quad \quad} & M_p(G) = \epsilon_p \text{Hom}(\Lambda, G) \\ \Lambda_p \otimes_{\text{Cart}_p(k)} M & \xleftarrow{\quad \quad \quad} & M \end{array}$$

(4.5.1) Lemma *Let k be a commutative $\mathbb{Z}_{(p)}$ -algebra with 1. Let $\alpha_p : L_p \rightarrow M_p$ be a surjective homomorphism V -reduced $\text{Cart}(k)$ -module. Let K_p be the kernel of α_p . Let K be the kernel of*

$$\text{id} \otimes \alpha_p : \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} L_p \rightarrow \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} M_p$$

with the induced filtration.

- (i) *The $\text{Cart}_p(k)$ -module K_p is V -reduced.*
- (ii) *The pair $(K, \text{Fil}^\bullet K)$ is the V -reduced $\text{Cart}(k)$ -module which corresponds to K_p under Thm. 4.4.*
- (iii) *The sequence $0 \rightarrow V^i K / V^{i+1} M \rightarrow V^i L / V^{i+1} L \rightarrow V^i M / V^{i+1} M \rightarrow 0$ is short exact for every $i \geq 0$.*

PROOF. The argument of 3.3.3 works here as well.

(4.5.2) Corollary *Let M_p be a V -reduced $\text{Cart}(k)$ -module. Then there exists an exact sequence*

$$\dots \xrightarrow{\partial_{i+1}} L_i \xrightarrow{\partial_i} L_{i-1} \xrightarrow{\partial_{i-1}} \dots \xrightarrow{\partial_1} L_0 \xrightarrow{\partial_0} M_p \rightarrow 0$$

of $\text{Cart}_p(k)$ modules such that L_i is a free V -reduced $\text{Cart}_p(k)$ -module, and induces an exact sequence on each V -adically graded piece.

(4.5.3) Definition (1) A *torsion right $\text{Cart}_p(k)$ -module Q* is a right $\text{Cart}_p(k)$ -module such that for every $x \in Q$, there exists a natural number $n \geq 0$ such that $x \cdot V^n = 0$.

- (2) Let Q be a torsion right $\text{Cart}(k)$ -module. Let M be a V -reduced $\text{Cart}_p(k)$ -module. Let L_\bullet be a resolution of M by free V -reduced $\text{Cart}_p(k)$ -modules as in 4.5.2. Define $\overline{\text{Tor}}_i(Q, M)$, $i \geq 0$, by

$$\overline{\text{Tor}}_i(Q, M) = H_i(Q \otimes_{\text{Cart}_p(k)} L_\bullet).$$

Exercise (i) Show that the continuous torsion functors $\overline{\text{Tor}}_\bullet^{\text{Cart}(k)}(Q, M)$ are well-defined.

- (ii) Let Q be a torsion right $\text{Cart}_p(k)$ -module and let $\widehat{\text{Cart}_p(k)}^{(I)}$ be a free V -reduced $\text{Cart}_p(k)$ -module with basis indexed by a set I . Prove that $Q \otimes_{\text{Cart}_p(k)} \widehat{\text{Cart}_p(k)}^{(I)}$ is naturally isomorphic to $Q^{(I)}$, the direct sum of copies of the abelian group Q indexed by I .
- (iii) Show that for any V -reduced $\text{Cart}_p(k)$ -module M and any short exact sequence $0 \rightarrow Q_1 \rightarrow Q_2 \rightarrow Q_3 \rightarrow 0$ of torsion right $\text{Cart}_p(k)$ -modules, one has a long exact sequence consisting of the abelian groups $\overline{\text{Tor}}_i^{\text{Cart}_p(k)}(Q_j, M)$.
- (iv) Show that for any torsion right $\text{Cart}_p(k)$ -module Q and any short exact sequence $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ of V -reduced $\text{Cart}_p(k)$ -modules, one has a long exact sequence consisting of the abelian groups $\overline{\text{Tor}}_i^{\text{Cart}_p(k)}(Q, M_j)$.

(4.5.4) Remark Let Q be a torsion right $\text{Cart}(k)$ -module.

(i) The canonical maps

$$Q \otimes_{\text{Cart}(k)} (\text{Cart}(k)\epsilon_p) \rightarrow Q \overline{\otimes}_{\text{Cart}(k)} (\text{Cart}(k)\epsilon_p) \rightarrow Q \epsilon_p$$

are isomorphisms of torsion right $\text{Cart}_p(k)$ -modules; denote these canonically isomorphic right $\text{Cart}_p(k)$ -modules by Q_p .

(ii) The subset $Q \epsilon_p$ of Q , the image of right multiplication by ϵ_p on Q , consists of all elements $x \in Q$ such that $x \cdot V_n = 0$ for all $n \geq 2$ with $(n, p) = 1$.

(4.5.5) Proposition Let Q be a torsion right $\text{Cart}(k)$ -module, and let M be a V -reduced $\text{Cart}(k)$ -module. Let $Q_p = Q \overline{\otimes}_{\text{Cart}(k)} \text{Cart}(k)\epsilon_p$. Let $M_p = \epsilon_p M$ be the V -reduced $\text{Cart}_p(k)$ -module consisting of all p -typical elements in M .

(i) The canonical map

$$Q_p \otimes_{\text{Cart}_p(k)} M_p = (Q \overline{\otimes}_{\text{Cart}(k)} \text{Cart}(k)\epsilon_p) \otimes_{\text{Cart}_p(k)} M_p \longrightarrow Q \overline{\otimes}_{\text{Cart}(k)} M$$

is an isomorphism.

(ii) For all $i \geq 0$ the canonical map

$$\overline{\text{Tor}}_i^{\text{Cart}_p(k)}(Q_p, M_p) \longrightarrow \overline{\text{Tor}}_i^{\text{Cart}(k)}(Q, M)$$

is an isomorphism.

PROOF. Since Q is a torsion right $\text{Cart}(k)$ -module, the canonical map

$$Q \otimes_{\text{Cart}(k)} (\text{Cart}(k)\epsilon_p \otimes_{\text{Cart}_p(k)} \epsilon_p M) \rightarrow Q \overline{\otimes}_{\text{Cart}(k)} (\text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} \epsilon_p M)$$

is an isomorphism, and (i) follows from the associativity of tensor product. To prove (ii), let $L_\bullet \rightarrow M$ be a resolution of M by free V -reduced $\text{Cart}(k)$ -modules. Then $\epsilon_p L_\bullet \rightarrow \epsilon_p M = M_p$ is a resolution of M_p by free V -reduced $\text{Cart}_p(k)$ -modules. By (i) the natural map

$$Q_p \otimes_{\text{Cart}_p(k)} \epsilon_p L_\bullet \xrightarrow{\sim} Q \overline{\otimes}_{\text{Cart}(k)} M$$

is an isomorphism of chain complexes, and the statement (ii) follows. ■

(4.5.6) Theorem Let k be a commutative $\mathbb{Z}_{(p)}$ -algebra with 1. Let N be a nilpotent k -algebra. Let M_p be a V -reduced $\text{Cart}_p(k)$ -module. Let $M = \text{Cart}(k)\epsilon_p \widehat{\otimes} M_p$.

(i) The canonical map

$$\Lambda_p(N) \otimes_{\text{Cart}_p(k)} M_p \longrightarrow \Lambda(N) \overline{\otimes}_{\text{Cart}(k)} M$$

is an isomorphism.

(ii) Assume either that M_p is V -flat, or that N has a finite decreasing filtration

$$N = \text{Fil}^1 N \supseteq \text{Fil}^2 N \supseteq \cdots \supseteq \text{Fil}^s N = (0)$$

such that each $\text{Fil}^j N$ is an ideal of N , $(\text{Fil}^j N)^2 \subseteq \text{Fil}^{j+1} N$ and $\text{Fil}^j N / \text{Fil}^{j+1} N$ is a flat k -module for $j = 1, \dots, s-1$. Then

$$\overline{\text{Tor}}_i^{\text{Cart}_p(k)}(Q_p, M_p) \cong \overline{\text{Tor}}_i^{\text{Cart}(k)}(Q, M) = (0) \quad \forall i \geq 1.$$

PROOF. The statement (i) is a corollary of 4.5.5 (i). The statement (ii) follows from 4.5.5 (ii) and Thm. 3.3.5. ■

(4.5.7) PROOF OF THM. 4.5. Theorem 4.5 follows from Thm. 4.5.6, Thm. 4.4 and Thm. 3.3. ■

(4.6) **Theorem** Let k be a commutative $\mathbb{Z}_{(p)}$ -algebra with 1.

(i) Let M be a V -reduced $\text{Cart}_p(k)$ -module. Let $\{x_i \mid i \in I\}$ be a family of elements in M indexed by a set I such that M/VM is a free k -module with basis $\{\bar{x}_i \mid i \in I\}$, where \bar{x}_i denotes the image of x_i in M/VM . Then

$$M = \left\{ \sum_{\substack{m \geq 0 \\ i \in I}} V^m \langle a_{mi} \rangle x_i \mid \begin{array}{l} \text{(i) } a_{mi} \in k \quad \forall m \geq 0, \forall i \in I \\ \text{(ii) } \forall m \exists \text{ a finite subset } J_m \subset I \\ \text{s.t. } a_{mi} = 0 \text{ or if } i \notin J_m \end{array} \right\}$$

(ii) Notation as in (i) above. There exists uniquely determined elements $a_{mij} \in k$, with $(m, i, j) \in \mathbb{N} \times I \times I$ such that

$$F x_i = \sum_{\substack{m \in \mathbb{N} \\ j \in I}} V^m \langle a_{mij} \rangle x_j, \quad \forall i \in I \quad \forall m \in \mathbb{N},$$

and for each $m \geq 0$ and each $i \in I$, $a_{mij} = 0$ for all j outside a finite subset of I .

(iii) Let $\alpha_{mij} \in W_p(k)$ be a family of elements in $W_p(k)$ indexed by $\mathbb{N} \times I \times I$ such that for each $m \geq 0$ and each $i \in I$, there exists a finite subset $J_{mi} \subset I$ such that $\alpha_{mij} = 0$ for all $j \notin J_{mi}$. Then the $\text{Cart}_p(k)$ -module N defined by the short exact sequence $\text{Cart}_p(k)$ -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & L_1 = \widehat{\text{Cart}_p(k)}^{(I)} & \xrightarrow{\psi} & L_2 = \widehat{\text{Cart}_p(k)}^{(I)} & \longrightarrow & N \longrightarrow 0 \\ & & f_i \longmapsto & & F e_i - \sum_{m,j} V^m \alpha_{mij} e_j & & \\ & & & & e_i \longmapsto & & y_i \end{array}$$

is V -reduced. Here $\{f_i \mid i \in I\}$, $\{e_i \mid i \in I\}$ are the bases of the two free V -reduced $\text{Cart}_p(k)$ -modules L_1 , L_2 respectively. Moreover the image of the elements $\{y_i \mid i \in I\}$ in N/VN form a k -basis of N/VN .

PROOF. The statement (i) follows from the definition of V -reduced $\text{Cart}_p(k)$ -modules, and (ii) follows from (i).

To prove (iii), it suffices to show that the sequence in the displayed formula induces an exact sequence

$$0 \rightarrow L_1/VL_1 \xrightarrow{\bar{\psi}} L_2/VL_2 \rightarrow N/VN \rightarrow 0$$

of k -modules, and the elements $(\bar{y}_i)_{i \in I}$ form a k -basis of N/VN . Here we used the convention that \bar{y}_i denotes the image of y_i in N/VN ; the same convention will be used for L_1/VL_1 and L_2/VL_2 . Recall that a typical element $\omega_p(\underline{c}) \in W_p(k)$ is identified with the element $\sum_{m \geq 0} V^m \langle c_m \rangle F^m$ of $\text{Cart}_p(k)$. For $i, j \in I$, let $a_{ij} = w_0(\alpha_{0ij}) \in k$, so that $\alpha_{0ij} - \omega_p(a_{ij}, 0, 0, \dots) \in V(W_p(k))$. We know that

$$\begin{aligned} L_1/VL_1 &= \bigoplus_{n \in \mathbb{N}, i \in I} \overline{F^n f_i}, & L_2/VL_2 &= \bigoplus_{n \in \mathbb{N}, i \in I} \overline{F^n e_i} \\ \bar{\psi} \left(\sum_{n \geq 0, i \in I} b_{ni} \overline{F^n f_i} \right) &= \sum_{n \geq 0, j \in I} \left(b_{n-1, j} - \sum_{i \in I} b_{ni} a_{i,j}^{p^n} \right) \overline{F^n e_j}. \end{aligned}$$

The desired conclusion follows from an easy calculation. ■

(4.6.1) Remark In the situation of 4.6 (i), (ii), we have a short exact sequence of V -reduced $\text{Cart}_p(k)$ -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & L_1 = \widehat{\text{Cart}_p(k)}^{(I)} & \longrightarrow & L_2 = \widehat{\text{Cart}_p(k)}^{(I)} & \longrightarrow & M \longrightarrow 0 \\ & & f_i \longmapsto & & F e_i - \sum_{m,j} V^m \langle a_{mij} \rangle e_j & & \\ & & & & e_i \longmapsto & & x_i \end{array}$$

The family of equations $Fx_i = \sum_{m,j} V^m \langle a_{mij} \rangle x_j$, $i \in I$, are called the *structural equations* of M for the generators $\{x_i \mid i \in I\}$.

(4.7) Proposition *Let M_p be a V -reduced V -flat $\text{Cart}_p(k)$ -module. Let k' be a commutative k algebra with 1. Let M'_p be the V -adic completion of the left $\text{Cart}_p(k')$ -module $\text{Cart}_p(k') \otimes_{\text{Cart}_p(k)} M_p$.*

(i) *The $\text{Cart}_p(k')$ -module M' is V -reduced, and $k' \otimes_k (M/VM) \xrightarrow{\sim} M'/VM'$ as k' -modules.*

(ii) *For every nilpotent k' -algebra N , there is a canonical isomorphism*

$$\Lambda_p(N) \overline{\otimes}_{\text{Cart}_p(k)} M_p \xrightarrow{\sim} \Lambda_p(N) \overline{\otimes}_{\text{Cart}_p(k')} M'_p.$$

(iii) *Suppose that $\{x_i \mid i \in I\}$ is a family of elements in M_p such that $\{\bar{x}_i \mid i \in I\}$ form a k -basis of M/VM . Let $Fx_i = \sum_{m \geq 0, j \in I} V^m \langle a_{mij} \rangle x_j$. $i \in I$ be the structural equation of M w.r.t. the generators $\{x_i \mid i \in I\}$. Then these equations are also the structural equations of M' for the generators $\{1 \otimes x_i \mid i \in I\}$.*

PROOF. Exercise. ■

(4.8) EXERCISES.

(4.8.1) Exercise Prove that the equivalence of categories in Thm. 4.5 extends to an equivalence of categories between the category of V -reduced V -flat $\text{Cart}_p(k)$ -modules and the category of functors $G : \mathfrak{Nilp}_k \rightarrow \mathfrak{Ab}$ which are right exact and weakly left exact.

(4.8.2) Exercise Prove that the left ideal $\text{Cart}(k)\epsilon_p$ of $\text{Cart}(k)$ consists of all elements $x \in \text{Cart}(k)$ such that $xV_n = 0$ for all $n \geq 2$ with $(n, p) = 1$. (Hint: Prove that $x \cdot (1 - \epsilon_p) \in \text{Fil}^m \text{Cart}(k) = 0$ for all $m \geq 1$. Or, use Exer. 2.7.2.)

(4.8.3) Exercise Let x be an element of $\text{Cart}(k)$.

- (i) Prove that $x \cdot \epsilon_p = 0$ if and only if x lies in the closure of the sum of left ideals $\sum_{(n,p)=1} \text{Cart}(k)F_n$. (Hint: Use 2.7.2.)
- (ii) Prove that $\epsilon_p \cdot x = 0$ if and only if x lies in the convergent sum of right ideals $\sum_{(n,p)=1} V_n \text{Cart}(k)$.

(4.8.4) Exercise Prove that

$$\epsilon_p \text{Cart}(k) = \left\{ \sum_{\substack{i,j \geq 0 \\ (n,p)=1}} V^i \langle a_{i,j,n} \rangle F^j F_n \mid \begin{array}{l} \text{(i) } a_{i,j,n} \in k \ \forall i, j \geq 0, \forall n \geq 1 \text{ with } (n, p) = 1, \\ \text{(ii) } \forall i \geq 0, \exists C_i > 0 \text{ s.t. } a_{i,j,n} = 0 \text{ if } j > C_i \text{ or } \\ n > C_i. \end{array} \right\}$$

In other words, $\epsilon_p \text{Cart}(k)$ is the V -adic completion of the discrete direct sum of the free $\text{Cart}_p(k)$ -modules $\text{Cart}_p(k) \cdot F_n$, where n ranges through all positive integers prime to p .

(4.8.5) Exercise Show that the canonical maps

$$\begin{aligned} \text{Cart}(k)\epsilon_p \widehat{\otimes}_{\text{Cart}_p(k)} \epsilon_p \text{Cart}(k) &\longrightarrow \text{Cart}(k) \\ \epsilon_p \text{Cart}(k) \otimes_{\text{Cart}_p(k)} \epsilon_p \text{Cart}(k) &\longrightarrow \text{Cart}_p(k) \end{aligned}$$

are isomorphisms.

(4.8.6) Exercise Let S be a subset of the set of all prime numbers, and let $\mathbb{Z}_S = \mathbb{Z}[\frac{1}{\ell}]_{\ell \notin S}$ be the subring of \mathbb{Q} generated by \mathbb{Z} and all prime numbers $\ell \notin S$. Generalize the results in this section to the case when the base ring k is a commutative algebra over \mathbb{Z}_S with 1.

(4.8.7) Exercise Assume that k is a field of characteristic p . Let M be a V -reduced $\text{Cart}_p(k)$ -module such that $\dim_k(M/VM) = 1$. Let $e \in M$ be an element of M such that $e \notin VM$, so that $M = \text{Cart}_p(k) \cdot e$. Suppose that $Fe = \sum_{m \geq n} V^m \langle a_m \rangle e$, with $a_m \in k$ for all $m \geq n$, and $a_n \neq 0$.

- (i) Suppose that there exists an element $b \in k$ such that $b^{p^{n+1}-1} = a_n$. Prove that there exists a generator x of M such that $Fx - V^n x \in V^{n+1}M$. (Hint: Use a generator of the form $\langle c \rangle e$.)
- (ii) Assume that k is perfect and there exists an element $b \in k$ such that $b^{p^{n+1}-1} = a_n$. Prove that there exists a generator y of M such that $Fy = V^n y$.

(4.8.8) Exercise Let k be a field of characteristic p . For $i = 1, 2$, let

$$M_i = \text{Cart}_p(k)/\text{Cart}_p(k) \cdot (F - \sum_{m \geq n_i} V^m \langle a_{im} \rangle),$$

where $a_{im} \in k$ for all $m \geq n_i$, and n_1, n_2 are natural numbers. If $n_1 \neq n_2$, prove that M_1 and M_2 are not isomorphic.

(4.8.9) Exercise Let $r \geq 1$ be a positive integer, and let $q = p^r$. Define formal power series $f(X) \in \mathbb{Q}[[X]]$ and $g(X, Y) \in \mathbb{Q}[[X, Y]]$ by

$$\begin{aligned} f(X) &= \sum_{n \geq 0} \frac{X^{q^n}}{p^n} \\ g(X, Y) &= f^{-1}(f(X) + f(Y)). \end{aligned}$$

It is well-known that $g(X, Y) \in \mathbb{Z}_{(p)}[[X, Y]]$ is a one-dimensional formal group law, a special case of the Lubin-Tate formal group law. The formal group law $g(X, Y)$ defines a smooth commutative formal group $G : \mathfrak{Nilp}_{\mathbb{Z}_{(p)}} \rightarrow \mathfrak{Ab}$. By definition, the $\text{Cart}(\mathbb{Z}_{(p)})$ -module M attached to G is $G(\mathbb{Z}_{(p)}[[X]]^+) = \mathbb{Z}_{(p)}[[X]]^+$. Let γ be the element of M corresponding to $X \in \mathbb{Z}_{(p)}[[X]]^+$

- (i) Prove that γ is a p -typical element of M . (Hint: Change the base ring from $\mathbb{Z}_{(p)}$ to \mathbb{Q} .)
- (ii) Prove that $M_p := \epsilon_p M$ is generated by γ .
- (iii) Prove that $F \cdot \gamma = V^{r-1} \cdot \gamma$.
- (iv) Prove that $\text{End}(G) = \mathbb{Z}_{(p)}$.

(4.8.10) Exercise Let k be a perfect field of characteristic p . Let n be a natural number. Prove that $\text{Cart}_p(k)/\text{Cart}_p(k) \cdot (F - V^n)$ is a free $W_p(k)$ -module of rank $n + 1$.

(4.8.11) Exercise Let $n \geq 0$ be a natural number. Let k be a field of characteristic p . Let $M = \text{Cart}_p(k)/\text{Cart}_p(k) \cdot (F - V^n)$.

- (i) If $n = 0$, show that M is a free $W_p(k)$ -module of rank one.
- (ii) Suppose that $n \geq 1$, $c \in k$. Prove that $(V^j \langle c \rangle F^j) \cdot M \subseteq M$ if $j \geq 2$.

(iii) Prove that M is *not* a free $W_p(k)$ -module if $n \geq 1$ and k is not perfect.

(4.8.12) Exercise Notation as in 4.8.11. Let k_1 be the finite subfield of k consisting of all elements $x \in k$ such that $x^{p^{n+1}} = x$. Let $\text{Card}(k_1) = p^r$.

(i) Show that $r|n+1$.

(ii) Show that $\text{End}_{\text{Cart}(k)}(M)$ is a $W_p(k_1)$ -module of rank $(n+1)$.

(iii) Let $D = \text{End}_{\text{Cart}(k)}(M) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$. Prove that D_p is a division algebra.

(iv) Prove that the center of D is a totally ramified extension of degree $e = \frac{n+1}{r}$, isomorphic to $\mathbb{Q}_p[T]/(T^e - p)$.

(v) Find the Brauer invariant of the division algebra D with center E .

§A. Appendix: Witt vectors

In this appendix we explain the basic properties of the ring \widetilde{W} of universal Witt vectors and the ring W_p of p -adic Witt vectors. Both are ring schemes over \mathbb{Z} , and W_p is a factor of \widetilde{W} over \mathbb{Z}_p . This appendix is logically independent of the rest of this article, and the main results in this set of notes does not depend on the Witt vectors.

(A.1) Definition The universal Witt vector group \widetilde{W} is defined as the functor from the category of all commutative algebras with 1 to the category of abelian groups such that

$$\widetilde{W}(R) = 1 + T R[[T]] \subset R[[T]]^\times$$

for every commutative ring R with 1. It turns out that the \widetilde{W} has a natural structure as a ring scheme. When we regard a formal power series $1 + \sum_{m \geq 1} u_m T^m$ in $R[[T]]$ as an element of $\widetilde{W}(R)$, we use the notation $\omega(1 + \sum_{m \geq 1} u_m T^m)$. It is easy to see that every element of $\widetilde{W}(R)$ has a unique expression as

$$\omega \left(\prod_{m \geq 1} (1 - a_m T^m) \right).$$

Hence \widetilde{W} is isomorphic to $\text{Spec } \mathbb{Z}[x_1, x_2, x_3, \dots]$ as a scheme; the R -valued point such that $x_i \mapsto a_i$ is denoted by $\omega(\underline{a})$, where \underline{a} is short for (a_1, a_2, a_3, \dots) . In other words, $\omega(\underline{a}) = \omega(\prod_{m \geq 1} (1 - a_m T^m))$

(A.1.1) Definition The ring structure of \widetilde{W} is given by

$$\omega(1 - a T^m) \cdot \omega(1 - b T^n) = \omega \left((1 - a^{\frac{n}{r}} b^{\frac{m}{r}} T^{\frac{mn}{r}})^r \right), \quad \text{where } r = (m, n).$$

(A.1.2) Exercise (i) Prove that the recipe above for multiplication indeed defines a ring structure on \widetilde{W} . In other words, prove that $(\omega(1-a_1 T^{n_1}) \cdot \omega(1-a_2 T^{n_2})) \cdot \omega(1-a_3 T^{n_3}) = \omega(1-a_1 T^{n_1}) \cdot (\omega(1-a_2 T^{n_2}) \cdot \omega(1-a_3 T^{n_3}))$ for any elements $a_1, a_2, a_3 \in R$ and any $n_1, n_2, n_3 \geq 1$.

(ii) Show that ring structure on \widetilde{W} is uniquely determined by the requirement that $\omega(1-aT) \cdot \omega(1-bT) = \omega(1-abT)$ for every commutative ring R and all elements $a, b \in R$.

(A.1.3) The group scheme \widetilde{W} has a decreasing filtration $(\text{Fil}^n \widetilde{W})_{n \geq 1}$, where

$$\text{Fil}^n \widetilde{W}(R) = \omega(1 + T^n R[[T]]) \subset \widetilde{W}(R).$$

In terms of the coordinates in §A.1, $\text{Fil}^n \widetilde{W}(R)$ consists of all R -valued points such that the coordinates x_1, \dots, x_{n-1} vanish. For every n , $\text{Fil}^n \widetilde{W}(R)$ is an ideal of $\widetilde{W}(R)$. This $W(R)$ is complete with respect to this filtration, and the addition, multiplication, and the operator F_n, V_n defined in A.3 below are continuous with respect to this filtration; see (A.3.1) (9).

(A.2) Definition There is a homomorphism of ring schemes

$$\text{ghost} : \widetilde{W} \longrightarrow \prod_{m=1}^{\infty} \mathbb{A}^1 = \text{Spec } k[\tilde{w}_1, \tilde{w}_2, \tilde{w}_3, \dots],$$

where the target has the standard ring scheme structure. The coordinates of the map ghost are

$$\tilde{w}_m(\underline{a}) = \sum_{d|m} d \cdot a_d^{m/d}.$$

Equivalently if we identify an R -valued point \underline{r} of $\text{Spec } k[\tilde{w}_1, \tilde{w}_2, \tilde{w}_3, \dots]$ with the power series $1 + \sum_{m=1}^{\infty} r_m T^m \in R[[T]]$ for a commutative ring R , then the map ghost on \widetilde{W} is induced by the operator $-t \frac{d}{dt} \log$:

$$\text{ghost}(\omega\left(\prod_{m \geq 1} (1 - a_m T^m)\right)) = \sum_{m \geq 1} \sum_{d \geq 1} m a_m^d T^{md} = \sum_{m \geq 1} \tilde{w}_m(\underline{a}) T^m.$$

(A.2.1) Exercise Prove that the map ghost is a homomorphism of ring schemes over \mathbb{Z} , and is an isomorphism over \mathbb{Q} .

(A.3) Definition There are two families of endomorphisms of the group scheme \widetilde{W} : V_n and F_n , $n \in \mathbb{N}_{\geq 1}$. Also for each commutative ring R with 1 and each element $c \in R$ we have an endomorphism $[c]$ of $\widetilde{W} \times_{\text{Spec } \mathbb{Z}} \text{Spec } R$. They are defined as follows

$$V_n : \omega(f(T)) \mapsto \omega(f(T^n))$$

$$F_n : \omega(f(T)) \mapsto \sum_{\zeta \in \mu_n} \omega(f(\zeta T^{\frac{1}{n}})) \quad (\text{formally})$$

$$[c] : \omega(f(T)) \mapsto \omega(f(cT))$$

The formula for $F_n(\omega(f(T)))$ means that $F_n(\omega(f(T)))$ is defined as the unique element such that $V_n(F_n(\omega(f(T)))) = \sum_{\zeta \in \mu_n} \omega(f(\zeta T))$.

(A.3.1) Exercise Prove the following statements.

- (1) $V_n(\omega(1 - aT^m)) = \omega(1 - aT^{mn})$, $F_n(\omega(1 - aT^m)) = \omega((1 - a^{\frac{n}{r}} T^{\frac{m}{r}})^r)$, $r = (m, n)$.
- (2) $F_n F_m = F_{mn}$, $V_m V_n = V_{mn}$.
- (3) $V_n F_m = F_m V_n$ if $(m, n) = 1$.
- (4) $F_n V_n = n$, i.e. $F_n V_n \omega(1 - a_m T^m) = \omega((1 - a_m T^m)^n)$ for all $m, n \geq 1$.
- (5) Let p be a prime number. Then $V_p F_p = p$ on $\widetilde{W}(R)$ if $p = 0$ in R . Conversely if $V_p(1) = V_p(F_p(1)) = p$, then $p = 0$ in R . (Hint: For the “only if” part, show that $V_p F_p \omega(1 - T) = \omega(1 - T^p)$.)
- (6) F_n is a ring homomorphism on \widetilde{W} for all $n \geq 1$. (Hint: Either verify this statement for the set of topological generators $\omega(1 - a_m T^m)$, or use the ghost coordinates.)
- (7) $x \cdot (V_n y) = V_n(F_n(x) \cdot y)$ for all $x, y \in \widetilde{W}(R)$.
- (8) If a positive integer N is invertible in R , then N is also invertible in $W(R)$.
- (9) $V_n(\text{Fil}^m \widetilde{W}) \subseteq \text{Fil}^{mn} \widetilde{W}$, $F_n(\text{Fil}^m \widetilde{W}) \subseteq \text{Fil}^{\lceil \frac{m}{n} \rceil} \widetilde{W}$, $\text{Fil}^m \widetilde{W} \cdot \text{Fil}^n \widetilde{W} \subseteq \text{Fil}^{\max(m, n)} \widetilde{W}$.
- (10) For all $c \in R$, all $m, n \geq 1$ and all $x \in \widetilde{W}(R)$, we have

$$\tilde{w}_m([c](x)) = c^m \tilde{w}_m(x), \quad \tilde{w}_m(F_n(x)) = \tilde{w}_{mn}(x), \quad \text{and} \quad \tilde{w}_m(V_n(x)) = \begin{cases} n \tilde{w}_{\frac{m}{n}} & \text{if } n|m \\ 0 & \text{if } n \nmid m \end{cases}$$

(A.3.2) Exercise Let R be a commutative ring with 1.

- (i) Show that every endomorphism Φ of the group scheme \widetilde{W} over R is determined by the element $\Phi_{R[X]}(1 - XT) \in \widetilde{W}(R[X])$.
- (ii) Prove that every element $\Phi \in \text{End}_R(\widetilde{W})$ can be expressed as an infinite series in the form

$$\sum_{m, n \geq 1} V_m [a_{mn}] F_n$$

with $a_{mn} \in R$ for all $m, n \geq 1$, and for every m there exists $C_m \geq 0$ such that $a_{mn} = 0$ if $n \geq C_m$. The elements $a_{mn} \in R$ are uniquely determined by the endomorphism Φ , and every family of elements $\{a_{mn}\}$ in R satisfying the above condition gives an endomorphism of \widetilde{W} .

(A.4) Definition Over $\mathbb{Z}_{(p)}$ we define a projector

$$\epsilon_p := \sum_{(n,p)=1} \frac{\mu(n)}{n} V_n F_n = \prod_{\ell \neq p} \left(1 - \frac{1}{\ell} V_\ell F_\ell\right),$$

where ℓ runs through all prime numbers not equal to p . Note that the factors $(1 - \frac{1}{\ell} V_\ell F_\ell)$ commute.

(A.4.1) Exercise Prove that

- (i) $(1 - \frac{1}{\ell} V_\ell F_\ell) V_\ell = 0 = F_\ell (1 - \frac{1}{\ell} V_\ell F_\ell)$. (Use A.3.1 (4).)
- (ii) $(1 - \frac{1}{\ell} V_\ell F_\ell)^2 = 1 - \frac{1}{\ell} V_\ell F_\ell$.
- (iii) $\epsilon_p \circ V_\ell = 0 = F_\ell \circ \epsilon_p \quad \forall \ell \neq p$.
- (iv) $\epsilon_p^2 = \epsilon_p$.

(A.4.2) Definition Denote by W_p the image of ϵ_p , i.e. $W_p(R) := \epsilon_p(\widetilde{W}(R))$ for every $\mathbb{Z}_{(p)}$ -algebra R . The functor W_p has a natural structure as a ring-valued functor induced from that of \widetilde{W} ; see Exer. A.4.5 (3); it is represented by the scheme $\text{Spec } \mathbb{Z}_{(p)}[y_0, y_1, y_2, \dots, y_n, \dots]$ according to the computation below.

(A.4.3) For every $\mathbb{Z}_{(p)}$ -algebra R and every sequence of elements $a_m \in R$, we have

$$\epsilon_p \left(\omega \left(\prod_{m \geq 1} (1 - a_m T^m) \right) \right) = \epsilon_p \left(\omega \left(\prod_{n \geq 0} (1 - a_{p^n} T^{p^n}) \right) \right) = \omega \left(\prod_{n \geq 0} E(a_{p^n} T^{p^n}) \right),$$

where

$$E(X) = \prod_{(n,p)=1} (1 - X^n)^{\frac{\mu(n)}{n}} = \exp \left(- \sum_{n \geq 0} \frac{X^{p^n}}{p^n} \right) \in 1 + X \mathbb{Z}_{(p)}[[X]]$$

is the inverse of the classical Artin-Hasse exponential. It follows that the map

$$\prod_0^\infty R \ni (c_0, c_1, c_2, \dots) \mapsto \omega \left(\prod_{n=0}^\infty E(c_n T^{p^n}) \right) \in W_p(R) = \epsilon_p(\widetilde{W}(R))$$

establishes a bijection between $\prod_0^\infty R$ and $W_p(R)$. Denote the element $\omega(\prod_{n=0}^\infty E(c_n T^{p^n})) \in W_p(R)$ by $\omega_p(\underline{c})$. We have shown that the functor $R \mapsto W_p(R)$ is represented by the scheme $\text{Spec } k[y_0, y_1, y_2, \dots]$, such that the element $\omega_p(\underline{c})$ has coordinates $\underline{c} = (c_0, c_1, c_2, \dots)$.

(A.4.4) The ghost coordinates on \widetilde{W} simplifies greatly when restricted to W_p . Most of them vanish: $\tilde{w}_m(\omega_p(\underline{c})) = 0$ if m is not a power of p for all \underline{c} . Let $w_n(\underline{c}) = \tilde{w}_{p^n}(\omega_p(\underline{c}))$ for all $n \geq 0$. Then

$$w_n(\underline{c}) = \sum_{i=0}^n p^{n-i} c_{n-i}^{p^i},$$

and $\text{ghost}(\omega_p(\underline{c})) = \sum_{n=0}^{\infty} w_n(\underline{c}) T^{p^n}$. For each n the map $\omega_p(\underline{c}) \mapsto (w_n(\underline{c}))_n$ is a homomorphism of ring schemes $W_p \rightarrow \prod_0^{\infty} \mathbb{A}^1$. The endomorphism V_p, F_p of the group scheme \widetilde{W} induces endomorphisms V, F of the group scheme W_p . Clearly $V(\omega_p(c_0, c_1, c_2, \dots)) = \omega_p(0, c_1, c_2, \dots)$ for all \underline{c} .

(A.4.5) **Exercise** Verify the following statements.

- (1) $\epsilon_p(1) \cdot x = \epsilon_p(x)$ for all $x \in \widetilde{W}(R)$. (Hint: Use A.3.1 (7).)
- (2) $\epsilon_p(1) = \omega(E(T))$, $\epsilon_p(1) \cdot \epsilon_p(1) = \epsilon_p(1)$
- (3) $\epsilon_p(x \cdot y) = \epsilon_p(x) \cdot \epsilon_p(y)$ for all $x, y \in \widetilde{W}(R)$. Hence $W_p(R)$ is a subring of $\widetilde{W}(R)$ whose unit element is $\epsilon_p(1)$.
- (4) $FV = p$ on $W_p(R)$.
- (5) $VF = p$ on $W_p(R)$ if $p = 0$ in R . Conversely if $V(1) = V(F(1)) = p$ then $p = 0$ in R . (Hint: For the “only if” part, show that $VF(1) = \omega(E(T^p))$, while $p = \omega(E(T)^p)$.)
- (6) $V(Fx \cdot y) = x \cdot Vy$ for all $x, y \in W_p(R)$.
- (7) $F(xy) = F(x) \cdot F(y)$ for all $x, y \in W_p(R)$.
- (8) $F(\omega_p(c_0, c_1, c_2, \dots)) = \omega_p(c_0^p, c_1^p, c_2^p, \dots)$ if $p = 0$ in R and $c_i \in R$ for all R .
- (9) For each $a \in R$, let $\langle a \rangle := \omega_p(a, 0, 0, 0, \dots) = \omega(E(cT))$. Then

$$\langle a \rangle \cdot \omega_p(\underline{c}) = \omega_p(ac_0, a^p c_1, a^{p^2} c_2, \dots)$$

for all \underline{c} .

- (10) $w_n \circ F = w_{n+1}$ for all $n \geq 0$, and

$$w_n \circ V = \begin{cases} p w_{n-1} & \text{if } n \geq 1 \\ 0 & \text{if } n = 0 \end{cases}$$

- (11) $F\langle a \rangle = \langle a^p \rangle$ for any $a \in R$.

(A.4.6) **Exercise** The group scheme W has a decreasing filtration $\text{Fil}^n W$, $n \geq 0$ defined by $\text{Fil}^n W = V^n W(R)$, that is $\text{Fil}^n W(R)$ consists of all elements of the form $\omega_p(\underline{c})$ such that $c_i = 0$ for all $i < n$. Verify the following properties of this filtration.

(i) For each commutative ring R over $\mathbb{Z}_{(p)}$, the ring $W_p(R)$ is complete with respect to the filtration $\text{Fil}^\bullet W_p(R)$.

(ii) For each $n \geq 0$, $\text{Fil}^n W_p(R)$ is an ideal of $W_p(R)$.

(iii) $V(\text{Fil}^n W_p(R)) \subseteq \text{Fil}^{n+1} W_p(R)$ for all $n \geq 0$.

(iv) $F(\text{Fil}^n W_p(R)) \subseteq \text{Fil}^{n-1} W_p(R)$ for all $n \geq 0$.

(A.4.7) Exercise Show that the universal polynomials defining the ring law for W_p all have coefficients in \mathbb{Z} , therefore the ring scheme W_p over $\mathbb{Z}_{(p)}$ has a canonical extension to \mathbb{Z} .

(A.4.8) Exercise Suppose that $p = 0$ in R . Prove that the ideal $V W_p(R)$ is generated by p if and only if R is perfect; i.e. the Frobenius map $x \mapsto x^p$ for R is surjective.

(A.4.9) Exercise Suppose that k is a perfect field of characteristic p . Prove that $W(k)$ is a complete discrete valuation ring with maximal ideal $V W(k) = p W(k)$ and residue field k .

(A.5) Ramified Witt vectors

Let \mathcal{O} is a complete discrete valuation ring such that the residue field is a finite field with q elements. Let π be a uniformizing element of \mathcal{O} . The ring scheme of ramified Witt vectors W_π is similar to the p -adic Witt vectors, but the formal completion of \mathbb{G}_m is replaced by the Lubin-Tate formal group. More precisely, the role of the logarithm of the Artin-Hasse exponential is played by the power series $f_\pi(X) := \sum_{n \geq 0} \frac{X^{q^n}}{\pi^n}$ in (1.5.2). Let $E_\pi(X) = f_\pi^{-1}(X)$, the inverse of $f_\pi(X)$; $E_\pi(X)$ has coefficients in \mathcal{O} . One can show that there exist polynomials $g_i(\underline{u}, \underline{v})$, $i = 0, 1, 2, \dots$, where $\underline{u} = (u_0, u_1, u_2, \dots)$, $\underline{v} = (v_0, v_1, v_2, \dots)$, such that

$$\sum_{m \geq 0}^{\Phi_\pi} E_\pi(u_m T^{q^m}) + \sum_{m \geq 0}^{\Phi_\pi} E_\pi(v_m T^{q^m}) = \sum_{i \geq 0}^{\Phi_\pi} E_\pi(g_i(\underline{u}, \underline{v})) T^{q^i}.$$

The above family of polynomials $g_i(\underline{u}, \underline{v})$ defines a group law on \mathbb{A}^∞ , denoted by W_π , called the ramified Witt vectors for (\mathcal{O}, π) . The phantom coordinates are

$$w_{\pi, n}(\underline{u}) := \sum_{i=0}^n \pi^{n-i} u_{n-i}^{q^i}, \quad n \geq 0.$$

Each $w_{\pi, n}$ defines a group homomorphism from W_π to \mathbb{G}_a . Moreover there is a canonical ring scheme structure on W_π such that each $w_{\pi, n}$ is a ring homomorphism from W_π to \mathbb{A}^1 .

References

- [R] M. Raynaud. “ p -torsion” du schéma de Picard. *Astérisque* 64:87–148, 1973.
- [H] M. Hazewinkel. *Formal Groups and Applications*. Academic Press, 1978.
- [L] M. Lazard. *Commutative Formal Groups*. LNM 443, Springer, 1975.
- [Z] T. Zink. *Cartiertheorie kommutativer former Gruppen*. Teubner, 1984