

Homework Set 5

DUE: Thurs. Oct. 19, 2006. Late papers accepted until 1:00 Friday.

Math 508, Fall 2006

Jerry Kazdan

1. Let V and W be normed linear spaces and $L : V \rightarrow W$ a linear map, so $L(X + Y) = LX + LY$ and $L(cX) = cLX$. Define the *norm* of L by

$$\|L\| := \sup_{X \neq 0} \frac{\|LX\|_W}{\|X\|_V}.$$

We say that L is *bounded* if $\|L\| < \infty$

- a) The set $\mathcal{L}(V, W)$ of all linear maps from V to W is itself a linear space — since one can add maps, $L + M$, and multiply them by scalars, cL . Show that $\|L\|$ defines a norm on $\mathcal{L}(V, W)$, that is,
- $\|L\| \geq 0$, with $\|L\| = 0$ only if $L = 0$,
 - $\|cL\| = |c|\|L\|$ for any scalar c ,
 - $\|L + M\| \leq \|L\| + \|M\|$ (triangle inequality).
- b) Show that $\|L\| = \sup_{\|X\|_V=1} \|LX\|_W$.
- c) Let $A = \begin{pmatrix} 0 & 1 & -1 \\ 1 & 2 & 0 \end{pmatrix}$ define a linear map from \mathbb{R}^3 to \mathbb{R}^2 with the usual euclidean norms. Show that A is bounded. (You need not compute $\|A\|$, only get an upper bound for it).
- d) If $L : \mathbb{R}^k \rightarrow \mathbb{R}^n$ is given by any $n \times k$ matrix and both \mathbb{R}^k and \mathbb{R}^n have the Euclidean norm, show that L is bounded.
- e) Show that if $L : \ell_2 \rightarrow \ell_2$ is defined by

$$LX := (c_1x_1, c_2x_2, c_3x_3, \dots),$$

where c_j is a bounded sequence, then L is bounded.

- f) Show that if $L : \ell_2 \rightarrow \ell_2$ is defined by

$$LX := (x_1, 2x_2, \dots, nx_n, \dots)$$

is *not* a bounded linear map.

2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ have the property that $|f(x) - f(\hat{x})| < c|x - \hat{x}|$ for all real x, \hat{x} , and where c is a constant. Given any starting point x_0 , define $x_j, j = 1, 2, \dots$, recursively by the rule

$$x_{j+1} = f(x_j).$$

If $c < 1$, prove that the x_j converge to some real number.

3. a) Let A be a subset of the real numbers. Prove that the following statements are equivalent:
- A is closed.
 - Every sequence $x_n \in A$ such that $\sum_{n=1}^{\infty} \|x_{n+1} - x_n\| < \infty$ converges to a limit in A .
- b) Show this is also true for subsets A of the plane \mathbb{R}^2 with the usual euclidian norm.
4. Let ℓ_2 be the usual normed linear space space of infinite sequences $X = (x_1, x_2, \dots)$ with finite norm: $\|X\| := \sqrt{\sum_{j=1}^{\infty} |x_j|^2} < \infty$. Prove that ℓ_2 is *complete*. [SUGGESTION: See the similar proof that ℓ_1 is complete on the class web page.]