

**Problem Set 0: Rust Remover**

DUE: These problems will not be collected.

*You should already know the techniques to do these problems, although they may take some thinking.*

1. Show that for any positive integer  $n$ , the number  $2^{n+2} + 3^{2n+1}$  is divisible by 7.
2. Say you have  $k$  linear algebraic equations in  $n$  variables; in matrix form we write  $AX = Y$ . Give a proof or counterexample for each of the following.
  - a) If  $n = k$  there is always *at most one* solution.
  - b) If  $n > k$  you can *always* solve  $AX = Y$ .
  - c) If  $n > k$  the nullspace of  $A$  has dimension greater than zero.
  - d) If  $n < k$  then for *some*  $Y$  there is *no* solution of  $AX = Y$ .
  - e) If  $n < k$  the *only* solution of  $AX = 0$  is  $X = 0$ .
3. Let  $A$  and  $B$  be  $n \times n$  matrices with  $AB = 0$ . Give a proof or counterexample for each of the following.
  - a)  $BA = 0$
  - b) Either  $A = 0$  or  $B = 0$  (or both).
  - c) If  $\det A = -3$ , then  $B = 0$ .
  - d) If  $B$  is invertible then  $A = 0$ .
  - e) There is a vector  $V \neq 0$  such that  $BAV = 0$ .
4. Let  $A$  be a matrix, not necessarily square. Say  $\mathbf{V}$  and  $\mathbf{W}$  are particular solutions of the equations  $A\mathbf{V} = \mathbf{Y}_1$  and  $A\mathbf{W} = \mathbf{Y}_2$ , respectively, while  $\mathbf{Z} \neq 0$  is a solution of the homogeneous equation  $A\mathbf{Z} = 0$ . Answer the following in terms of  $\mathbf{V}$ ,  $\mathbf{W}$ , and  $\mathbf{Z}$ .
  - a) Find some solution of  $A\mathbf{X} = 3\mathbf{Y}_1$ .
  - b) Find some solution of  $A\mathbf{X} = -5\mathbf{Y}_2$ .
  - c) Find some solution of  $A\mathbf{X} = 3\mathbf{Y}_1 - 5\mathbf{Y}_2$ .
  - d) Find another solution (other than  $\mathbf{Z}$  and 0) of the homogeneous equation  $A\mathbf{X} = 0$ .
  - e) Find *two* solutions of  $A\mathbf{X} = \mathbf{Y}_1$ .

- f) Find another solution of  $A\mathbf{X} = 3\mathbf{Y}_1 - 5\mathbf{Y}_2$ .
- g) If  $A$  is a square matrix, then  $\det A = ?$
- h) If  $A$  is a square matrix, for any given vector  $\mathbf{W}$  can one always find at least one solution of  $A\mathbf{X} = \mathbf{W}$ ? Why?
5. a) If  $r (\neq 0)$  is a rational number and  $x$  is irrational, show that both  $r + x$  and  $rx$  are *irrational*.
- b) Prove that there is no rational number whose square is 12.
- c) Graph the points  $(x, y)$  in the plane  $\mathbb{R}^2$  that satisfy  $|y - x| > 2$ .
6. a) Write the complex number  $z = \frac{1}{a + ib}$  in the form  $c + id$ , where  $a, b, c$  and  $d$  are real numbers. Of course assume  $a + ib \neq 0$ .
- b) If  $w \in \mathbb{C}$  satisfies  $|w| = 1$ , show that  $1/w = \bar{w}$ . [ $\mathbb{C}$  is the set of complex numbers.]
7. Let  $z, w, v \in \mathbb{C}$  be complex numbers.
- a) Show that  $|z - w| \geq |z - v| - |v - w|$ .
- b) Graph the points  $z = x + iy$  in the complex plane that satisfy  $1 < |z - i| < 2$ .
- c) Let  $z, w \in \mathbb{C}$  be complex numbers with  $|z| < 1$  and  $|w| = 1$ . Show that

$$\left| \frac{w - z}{1 - \bar{z}w} \right| = 1.$$

8. a) Find a  $2 \times 2$  matrix that rotates the plane by  $+45$  degrees ( $+45$  degrees means 45 degrees *counterclockwise*).
- b) Find a  $2 \times 2$  matrix that rotates the plane by  $+45$  degrees followed by a reflection across the horizontal axis.
- c) Find a  $2 \times 2$  matrix that reflects across the horizontal axis followed by a rotation the plane by  $+45$  degrees.
- d) Find a matrix that rotates the plane through  $+60$  degrees, keeping the origin fixed.
- e) Find the inverse of each of these maps.
9. Let the continuous function  $f(\theta)$ ,  $0 \leq \theta \leq 2\pi$  represent the temperature along the equator at a certain moment, say measured from the longitude at Greenwich.. Show there are antipodal points with the *same* temperature.

10. A certain function  $f(x)$  has the property that  $\int_0^x f(t) dt = e^x \cos x + C$ . Find both  $f$  and the constant  $C$ .
11. If  $b \geq 0$ , show that for every real  $c$  the equation  $x^5 + bx + c = 0$  has exactly one real root.
12. Let  $p(x) := x^3 + cx + d$ , where  $c$ , and  $d$  are real. Under what conditions on  $c$  and  $d$  does this has three distinct real roots? [HINT: Sketch a graph of this cubic. Observe that if there are three distinct real roots then there is a local maximum and the polynomial is positive there. What about a local min?].
13. Prove that the function  $\sin x$  is not a polynomial. That is, there is no polynomial

$$p(x) = a_0 + a_1x + \dots + a_nx^n$$

with real coefficients so that  $\sin x = p(x)$  for all real numbers  $x$ . In your proof you can use any of the standard properties of the function  $\sin x$ .

[Last revised: May 11, 2014]