

Math 240: Linear Systems and Rank of a Matrix

Ryan Blair

University of Pennsylvania

Friday February 17, 2012

Outline

- 1 Systems of Linear Equations
- 2 linear Independence

Echelon Forms

Definition

A matrix is in **row-echelon form** if

- 1 Any row consisting of all zeros is at the bottom of the matrix.
- 2 For all non-zero rows the leading entry must be a one. This is called the **pivot**.
- 3 In consecutive rows the pivot in the lower row appears to the right of the pivot in the higher row.

Definition

A matrix is in **reduced row-echelon form** if it is in row-echelon form and every pivot is the only non-zero entry in its column.

Row Operations

We will be applying row operations to augmented matrices to find solutions to linear equations. This is called **Gaussian** or **Gauss-Jordan** elimination.

Here are the row operations:

- 1 Multiply a row by a number.
- 2 Switch rows.
- 3 Add a multiple of one row to another.

Row Operations

We will be applying row operations to augmented matrices to find solutions to linear equations. This is called **Gaussian** or **Gauss-Jordan** elimination.

Here are the row operations:

- 1 Multiply a row by a number.
- 2 Switch rows.
- 3 Add a multiple of one row to another.

Key Fact: If you alter an augmented matrix by row operations you preserve the set of solutions to the linear system.

Today's Goals

- 1 Be able to use rank of a matrix to determine if vectors are linearly independent.
- 2 Be able to use rank of an augmented matrix to determine consistency or inconsistency of a system.

Linear Independence

Definition

Let v_1, \dots, v_m be vectors in \mathbb{R}^n . The set $S = \{v_1, \dots, v_m\}$ is **linearly independent** if $c_1 v_1 + c_2 v_2 + \dots + c_n v_n = 0$ implies $c_1 = c_2 = \dots = c_n = 0$.

If there exists a non trivial solution to $c_1 v_1 + c_2 v_2 + \dots + c_n v_n = 0$ we say the set S is linearly dependant.

Linear Independence

Definition

Let v_1, \dots, v_m be vectors in \mathbb{R}^n . The set $S = \{v_1, \dots, v_m\}$ is **linearly independent** if $c_1 v_1 + c_2 v_2 + \dots + c_n v_n = 0$ implies $c_1 = c_2 = \dots = c_n = 0$.

If there exists a non trivial solution to $c_1 v_1 + c_2 v_2 + \dots + c_n v_n = 0$ we say the set S is linearly dependant.

Example: Are the following vectors linearly independent?

$$\langle 1, 2, 1 \rangle, \langle 1, 1, 0 \rangle, \langle 1, 0, 1 \rangle$$

Definition

Let A be an $m \times n$ matrix. The **rank** of A is the maximal number of linearly independent row vectors

Definition

(Pragmatic)

Let A be an $m \times n$ matrix and B be its row-echelon form. The **rank** of A is the number of pivots of B .

Definition

Let A be an $m \times n$ matrix. The **rank** of A is the maximal number of linearly independent row vectors

Definition

(Pragmatic)

Let A be an $m \times n$ matrix and B be its row-echelon form. The **rank** of A is the number of pivots of B .

Example What is the rank of the following matrix.

$$\begin{pmatrix} 2 & 0 & 1 & -1 \\ 0 & 1 & 2 & 1 \\ 2 & -1 & -1 & -2 \end{pmatrix}$$

Determining Linear independence Using Matrices

How to find if m vectors are linearly independent:

- 1 Make the vectors the rows of a $m \times n$ matrix (where the vectors are of size n)
- 2 Find the rank of the matrix.
- 3 If the rank is m then the vectors are linearly independent. If the rank is less than m , then the vectors are linearly dependent.

Determining Linear independence Using Matrices

How to find if m vectors are linearly independent:

- 1 Make the vectors the rows of a $m \times n$ matrix (where the vectors are of size n)
- 2 Find the rank of the matrix.
- 3 If the rank is m then the vectors are linearly independent. If the rank is less than m , then the vectors are linearly dependent.

Example: Are the following vectors linearly independent?

$$\langle -2, 0, 4, 1 \rangle, \langle 0, 0, 1, -1 \rangle, \langle 0, 1, 0, 1 \rangle, \langle 3, 2, -3, 0 \rangle$$

Determining Consistency

Given the linear system $Ax = B$ and the augmented matrix $(A|B)$.

- 1 If $\text{rank}(A) = \text{rank}(A|B) =$ the number of rows in x , then the system has a unique solution.
- 2 If $\text{rank}(A) = \text{rank}(A|B) <$ the number of rows in x , then the system has ∞ -many solutions.
- 3 If $\text{rank}(A) < \text{rank}(A|B)$, then the system is inconsistent.