

More Power Series Solutions to D.E.s at Regular Singular Points

Ryan Blair

University of Pennsylvania

Friday April 20, 2012

Outline

1 Review

Given a differential equation $y'' + P(x)y' + Q(x)y = 0$

Definition

A point x_0 is an **ordinary point** if both $P(x)$ and $Q(x)$ are analytic at x_0 . If a point is not ordinary it is a **singular point**.

Given a differential equation $y'' + P(x)y' + Q(x)y = 0$

Definition

A point x_0 is an **ordinary point** if both $P(x)$ and $Q(x)$ are analytic at x_0 . If a point is not ordinary it is a **singular point**.

Definition

A point x_0 is a **regular singular point** if the functions $(x - x_0)P(x)$ and $(x - x_0)^2Q(x)$ are both analytic at x_0 . Otherwise x_0 is irregular.

Theorem

(Frobenius' Theorem)

If x_0 is a regular singular point of $y'' + P(x)y' + Q(x)y = 0$, then there exists a solution of the form

$$y = \sum_{n=0}^{\infty} c_n (x - x_0)^{n+r}$$

where r is some constant to be determined and the power series converges on a non-empty open interval containing x_0

Theorem

(Frobenius' Theorem)

If x_0 is a regular singular point of $y'' + P(x)y' + Q(x)y = 0$, then there exists a solution of the form

$$y = \sum_{n=0}^{\infty} c_n (x - x_0)^{n+r}$$

where r is some constant to be determined and the power series converges on a non-empty open interval containing x_0

To solve $y'' + P(x)y' + Q(x)y = 0$ at a regular singular point x_0 , substitute

$$y = \sum_{n=0}^{\infty} c_n (x - x_0)^{n+r}$$

and solve for r and the c_n to find a series solution centered at x_0 . **We may not find two linearly independent solutions this way!**